Submitted to the Annals of Statistics

arXiv:1203.4401v2 [math.ST] 9 May 2012

MAXIMUM SMOOTHED LIKELIHOOD ESTIMATORS FOR THE INTERVAL CENSORING MODEL

By Piet Groeneboom

Delft University

We study the maximum smoothed likelihood estimator (MSLE) for interval censoring, case 2. Characterizations in terms of convex duality conditions are given and strong consistency is proved. Moreover, we show that, under smoothness conditions on the underlying distributions and using the usual bandwidth choice in density estimation, the local convergence rate is $n^{-2/5}$, in contrast with the rate $n^{-1/3}$ of the ordinary maximum likelihood estimator. In this situation the MSLE is asymptotically equivalent to the solution of a non-linear integral equation, which we solve using the implicit function theorem in Banach spaces. It is shown that, again under appropriate conditions, the MSLE has a normal limit distribution of which the expectation and variance are explicitly determined as a function of the underlying distributions. This is done by showing that the solution of the non-linear integral equation is sufficiently close to the solution of a linear integral equation, which, in turn, is sufficiently close to a "toy estimator", depending on the underlying distributions, for which we can compute the asymptotic bias and variance explicitly.

1. Introduction. In [8] the smoothed maximum likelihood estimator (SMLE) and maximum smoothed likelihood estimator (MSLE) were studied for the current status model, the simplest interval censoring model. It is called the interval censoring, case 1, model in [5] and [11]. It was shown in [8] that, under certain regularity conditions, the SMLE and the MSLE, evaluated at a fixed interior point, converge at rate $n^{-2/5}$ to the real underlying distribution function, if one takes a bandwidth of order $n^{-1/5}$. This convergence rate is faster than the convergence rate of the non-smoothed maximum likelihood estimator, which is $n^{-1/3}$ in this situation, as shown in [5] and [11]. Moreover, the limit distribution is normal, in contrast with the limit distribution of the non-smoothed maximum likelihood estimator.

The more realistic interval censoring model, where there is an interval in which the relevant (unobservable) event takes place, is much more common and also more important, in particular in medical statistics. It is called the interval censoring, case 2, model in [5] and [11]. A preliminary discussion of the SMLE in this situation can be found in [10], where it was shown that the development of the theory of the SMLE for this model crucially depends on a further analysis of the integral equations, studied in [2], [3] and [4]. In the present paper we study the MSLE instead and prove a consistency and asymptotic normality result for this estimator. We also discuss algorithms for computing the MSLE, which is a rather non-trivial issue.

The MSLE has the advantage over the SMLE that it can be used in situations where the SMLE cannot be used. As an example, the MLE itself is proved to be inconsistent for the current status continuous mark model (see [13]), and the SMLE will inherit the bad properties of the MLE in this situation, and also be inconsistent. On the other hand, a version of the MSLE, based on histograms, is proved to be consistent for this model in [9]. A similar phenomenon holds for the two-dimensional

---

AMS 2000 subject classifications: Primary 62G05, 62N01; secondary 62G20

Keywords and phrases: Interval censoring, maximum smoothed likelihood estimators, asymptotic distribution, integral equations, kernel estimators

right-censoring model, where the MLE is inconsistent (see [14]) and the SMLE will not make this better. In this case the MSLE will, under appropriate smoothing of the observation distribution, also be consistent.

We recall the interval censoring, case 2, model. Let $X_1,\dots,X_n$ be a sample of unobservable random variables from an unknown distribution function F_0 on $[0,\infty)$. Suppose that one can observe n pairs (T_i,U_i), independent of X_i, where $U_i>T_i$. Moreover,

$$\Delta_{i1}\stackrel{\text{def}}{=}1_{\{X_i\le T_i\}},\ \Delta_{i2}\stackrel{\text{def}}{=}1_{\{T_i<X_i\le U_i\}}\text{ and }\Delta_{i3}\stackrel{\text{def}}{=}1-\Delta_{i1}-\Delta_{i2},$$

provide the only information one has on the position of the random variables X_i with respect to the observation times T_i and U_i. In this set-up one wants to estimate the unknown distribution function F_0, generating the "unobservables" X_i.

Interestingly, from a computational point of view, the MLE for the distribution function of the hidden variable in the case that one has more observation times $T_i,U_i,V_i,\dots$ "per hidden variable", can always be reduced to the case of interval censoring, case 2. This follows from the fact that at most two of the observation times of the set $\{T_i,U_i,V_i,\dots\}$ are relevant for the location of the hidden variable. If we know that the hidden variable is located between two observation times, while the other observation times for this hidden variable are either more to the right or more to the left, then these other observation times do not give extra information and can be discarded in computing the MLE. Likewise, if we know that the hidden variable lies to the right of all these observation times, all observation times smaller than the largest one do not give extra information, with a similar situation if we know that the hidden observation time lies to the left of the smallest observation time for this variable. So in the last two cases only one observation time gives relevant information and the other ones can be discarded. This motivates concentrating on the interval censoring, case 2, model, as an extension of the current status model.

The MSLE for the interval censoring, case 2, model is defined in the following way. Let g be the joint density of the observation pairs (T_i,U_i), with first marginal g_1 and second marginal g_2. Moreover, let the densities h_{01}, h_{02} and h_0 be defined by

$$h_{01}(t)=F_0(t)g_1(t),\qquad h_{02}(u)=\{1-F_0(u)\}g_2(u),\qquad h_0(t,u)=\{F_0(u)-F_0(t)\}g(t,u).\tag{1.1}$$

We define $\tilde h_{nj}$, $j=1,2$, and $\tilde h_n$ as the estimates of the densities h_{0j}, $j=1,2$, and the 2-dimensional density h_0, where

$$\tilde h_{n1}(t)=\frac1n\sum_{i=1}^n K_{b_n}(t-T_i)\,\Delta_{i1},\qquad \tilde h_{n2}(u)=\frac1n\sum_{i=1}^n K_{b_n}(u-U_i)\Delta_{i3},\tag{1.2}$$

$$\tilde h_n(t,u)=\frac1n\sum_{i=1}^n K_{b_n}(t-T_i)\,K_{b_n}(u-U_i)\Delta_{i2},\tag{1.3}$$

and

$$K_{b_n}(x)=\frac{1}{b_n}K\left(\frac{x}{b_n}\right),$$

for a symmetric continuously differentiable kernel K with compact support, like the triweight kernel

$$K(x)=\tfrac{35}{32}\left(1-x^2\right)^3 1_{[-1,1]}(x),\ x\in\mathbb{R}.\tag{1.4}$$

At boundary points we use a boundary correction by replacing the kernel K by a linear combination of $K(u)$ and $uK(u)$. For example, if $t \in [0, b_n)$, we define

$$\tilde{h}_{n1}(t) = \alpha(t/b_n)\frac{1}{n}\sum_{i=1}^{n} K_{b_n}(t - T_i)\,\Delta_{i1} + \beta(t/b_n)\frac{1}{n}\sum_{i=1}^{n}\frac{t - T_i}{b_n}K_{b_n}(t - T_i)\,\Delta_{i1},$$

where the coefficients $\alpha(u)$ and $\beta(u)$ are defined by

$$\alpha(u)\int_{-1}^{u} K(x)\,dx + \beta(u)\int_{-1}^{u} xK(x)\,dx = 1,\ u \in [0, 1]. \tag{1.5}$$

and

$$\alpha(u)\int_{-1}^{u} xK(x)\,dx + \beta(u)\int_{-1}^{u} x^2K(x)\,dx = 0,\ u \in [0, 1]. \tag{1.6}$$

It may happen that $\tilde{h}_{n1}(t) < 0$; in that case we put $\tilde{h}_{n1}(t) = 0$.

If $t \in (M - b_n, M]$, we similarly define:

$$\tilde{h}_{n1}(t) = \alpha((M - t)/b_n)\frac{1}{n}\sum_{i=1}^{n} K_{b_n}(t - T_i)\,\Delta_{i1} - \beta((M - t)/b_n)\frac{1}{n}\sum_{i=1}^{n}\frac{t - T_i}{b_n}K_{b_n}(t - T_i)\,\Delta_{i1},$$

where the functions α and β are again defined by (1.5) and (1.6). The estimates $\tilde{h}_{n2}$ and $\tilde{h}_n$ are similarly defined if one or more (in the case of $\tilde{h}_n$) arguments have distance less than b_n to the boundary; for $\tilde{h}_n$ we apply this to the factors of the product of the kernels separately, in the same way as for the one-dimensional estimates $\tilde{h}_{nj}$.

The MSLE $\hat{F}_n$ is now defined as the distribution function, maximizing the criterion function

$$\ell(F) = \int \tilde{h}_{n1}(t)\log F(t)\,dt + \int \tilde{h}_{n2}(u)\{1 - F(t)\}\,du + \int \tilde{h}_n(t, u)\log\{F(u) - F(t)\}\,dt\,du, \tag{1.7}$$

as a function of the distribution function F. But in practice we discretize, and maximize:

$$\begin{aligned}\sum_{i=1}^{m}\bigl\{\tilde{h}_{n1}(t_i)\log F(t_i)\bigr\}d_i &+ \sum_{i=1}^{m}\bigl\{\tilde{h}_{n2}(t_i)\log\{1 - F(t_i)\}\bigr\}d_i\\ &+ \sum_{i=1}^{m-1}\sum_{j=i+1}^{m}\bigl\{\tilde{h}_n(t_i, t_j)\log\{F(t_j) - F(t_i)\}\bigr\}d_id_j,\end{aligned} \tag{1.8}$$

where $0 = t_0 < t_1, \dots < t_m$ are the points of a grid, and $d_i = t_i - t_{i-1}$, $i = 1, \dots, m$.

Note that the maximization of (1.7) is the same as the minimalization of the Kullback-Leibler distance

$$\begin{aligned}\int \tilde{h}_{n1}(t)\log\frac{\tilde{h}_{n1}(t)}{F(t)g_1(t)}\,dt &+ \int \tilde{h}_{n1}(t)\log\frac{\tilde{h}_{n2}(t)}{\{1 - F(t)\}g_2(t)}\,dt\\ &+ \int \tilde{h}_n(t, u)\log\frac{\tilde{h}_n(t, u)}{\{F(u) - F(t)\}g(t, u)}\,dt\,du\end{aligned}$$

Throughout this paper, we will assume that the following conditions is satisfied.

Condition 1.1. (Conditions on underlying distributions) *The observation density g is continuous and bounded on its support S, which is the closure of an open connected set, contained in $[0,M]^2$, and has the property*

$$\sup_{u-t<\varepsilon,\,0\le t<u\le M} g(t,u)=0,$$

for some $\varepsilon>0$. We call this the strict separation hypothesis. *Furthermore, we suppose that*

$$\{(t,u): g(t,u)>0\}$$

is a set bounded by the two line segments connecting $(0,0)$ and $(0,M)$, and $(0,M)$ and (M,M), respectively, and a smooth curve

$$\{(t,\gamma(t):t\in[0,M]\},$$

where $\varepsilon\le\gamma(0)<\gamma(M)=M$ and γ is an continuous function which is strictly increasing on the interval $[0,c]$, where c is defined by

$$c=\sup\{t>0:\gamma(t)<M\}.$$

The prototype of such a function is:

$$\gamma(t)=\begin{cases} t+\varepsilon, & t\in[0,M-\varepsilon),\\ M, & t\in[M-\varepsilon,M].\end{cases}$$

We will in fact assume that $(t,\gamma(t))$ runs inside the region bounded by two regions of this simple type for an $\varepsilon>0$ and $\varepsilon'>\varepsilon$, where $\varepsilon'<M/2$.

We also assume that the underlying distribution function F_0 of the "unobservables" is continuous on $(0,M)$ and that F_0 and the marginals g_1 and g_2 of g satisfy

$$\inf_{t\in(0,M)}\{g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t)\}\ge\delta, \tag{1.9}$$

for some $\delta>0$. Condition (1.9) was also used in [2], [3] and [4]. We denote the supports of g_1 and g_2 by S_1 and S_2, respectively, and assume that S_1 and S_2 are closed subintervals of $[0,M]$ such that $0\in S_1$ and $M\in S_2$.

Condition 1.2. (Conditions on the kernel estimators) *We assume that $\tilde h_{nj}$ and $\tilde h_n$ are kernel estimators of h_{0j} and h_0, respectively, defined by (1.2) and (1.3), respectively, for a symmetric continuously differentiable kernel K of type (1.4), with compact support. Moreover, for points near the boundary, boundary kernels are used, with coefficients $\alpha(t)$ and $\beta(t)$, defined by (1.5) and (1.6), respectively, where the functions α, β, and its derivatives α' and β' are bounded on $[0,1]$. Let a and b be defined by:*

$$a=\inf\left\{t\in[0,M]:\tilde h_{n1}(t)\vee\int_{u=0}^t \tilde h_n(u,t)\,du>0\right\}, \tag{1.10}$$

and

$$b=\sup\left\{t\in[0,M]:\tilde h_{n2}(t)\vee\int_{u=t}^M \tilde h_n(t,u)\,du>0\right\}, \tag{1.11}$$

We further assume that $0\le a<b\le M$ and that

$$\tilde h_{n1}(t)\vee\int_{u=0}^t \tilde h_n(u,t)\,du>0,\ t\in(a,M], \tag{1.12}$$

$$\tilde h_{n2}(t)\vee\int_{u=t}^M \tilde h_n(t,u)\,du>0,\ t\in[0,b). \tag{1.13}$$

An example of a kernel estimates, satisfying the conditions of Condition 1.2, is given by kernel estimates which use the triweight kernel, defined by (1.4). For this kernel, the weight functions α and β, used in constructing the boundary kernel, are decreasing on $[0,1]$, with $\alpha(0)\approx 6.11457$, $\alpha(1)=1$, and $\beta(0)\approx 15.0476$, $\beta(1)=0$. The corresponding derivatives α' and β' are increasing on $[0,1]$, with $\alpha'(0)\approx -40.8931$, $\alpha'(1)=0$, and $\beta'(0)\approx -100.635$, $\beta'(1)=0$.

Using Condition 1.1, we give a characterization in terms of necessary and sufficient (duality) conditions for the MSLE in section 2. In that section we also prove consistency of the MSLE, using techniques, similar to the method, used in [11], Part II, section 4.

In section 3 we discuss algorithms for computing the MSLE: the EM algorithm, an interior point algorithm and an iterative convex minorant algorithm. The iterative convex minorant algorithm is an adapted version of the algorithm, introduced in [5] and (again in) [11]. It turns out that the latter algorithm performs best in our experiments. The EM algorithm is very slow and therefore not suitable for larger sample sizes or simulation purposes.

In section 4 we will prove asymptotic normality of the MSLE at a fixed interior point of the domain of definition (Theorem 4.1). To give an example of a situation where the conditions of Theorem 4.1 are satisfied, we take g uniform on the triangle with vertices $(0,\varepsilon)$, $(0,1)$ and $(1-\varepsilon,1)$. This example is also discussed in [10]. Then

$$g_1(u)=\frac{2(1-u-\varepsilon)}{(1-\varepsilon)^2}1_{[0,1-\varepsilon]}(u)\,,\qquad g_2(v)=\frac{2(v-\varepsilon)}{(1-\varepsilon)^2}1_{[\varepsilon,1]}(v)\,, \tag{1.14}$$

and $g=2/(1-\varepsilon)^2$ on the triangle of its support. Pictures of the MSLE and MLE are given in Figures 1 and 2, where we took

$$F_0(x)=1-(1-x)^2,\ x\in[0,1].$$

Note that the density f_0 vanishes at the right boundary point, so, strictly speaking, the conditions of Theorem 4.1 are not satisfied (indicating that these conditions are probably slightly stronger than needed). The triweight kernel was used as smoothing kernel and the bandwidth was simply $b_n=n^{-1/5}$. Boundary kernels were used in computing $\tilde h_{n1}$ and $\tilde h_{n2}$, and for $\tilde h_n$ we just used the product of the kernels, again using a boundary correction using boundary kernels in each component, with bandwidth $b_n=n^{-1/5}$ for both coordinates. The separating gap ε was chosen to be $\varepsilon=0.1$. Figures 1 and 2 show the rather large improvement of the MSLE over the MLE when the underlying distribution is smooth.

In [8] an asymptotic normality result, analogous to Theorem 4.1 was proved for the current status model. In this case it is much easier to prove a result of this type, since we have an explicit characterization of the MSLE as the slope of the greatest convex minorant of a continuous cusum diagram. Moreover it was shown that the MSLE was asymptotically equivalent to a plug-in estimator, for which the asymptotic normality result could be easily proved. For the interval censoring, case 2, model, we have neither of these properties: the MSLE has to be computed by an iterative procedure, there is not an obvious characterization as a slope of a greatest convex minorant and there is no simple asymptotically equivalent plug-in estimator. Instead we have to rely on the implicit characterization, given in Lemma 2.1 of section 2. Furthermore, the role of the plug-in estimator is now played by the solution of a non-linear integral equation, see Lemmas 4.1 and 4.3. An implicit function theorem in Banach spaces is the basic tool for showing the existence and uniqueness of the solution to this equation, where the parameters are given by the estimates $\tilde h_{nj}$ and $\tilde h_n$ of h_{0j} and h_0, respectively.

In this paper we concentrate on the "separated case", where $U_i-T_i\ge\varepsilon$ for some $\varepsilon>0$, as in [2] and [3]. This case seems to be the most important case, and also to be the usual situation in



FIG 1. *The MSLE (solid) and MLE (dashed) on* $[0,1]$ *for a sample of size* $n = 100$ *from the distribution function* $F_0(x) = 1-(1-x)^2$ *(dotted);* g *is uniform on the triangle with vertices* $(0,\varepsilon)$, $(0,1)$ *and* $(1-\varepsilon,1)$, *where* $\varepsilon = 0.1$. *The bandwidth for the computation of the MSLE was* $b_n = n^{-1/5} \approx 0.398107$.



FIG 2. *The MSLE (solid) and MLE (dashed) on* $[0,1]$ *for a sample of size* $n = 1000$ *from the distribution function* $F_0(x) = 1-(1-x)^2$ *(dotted);* g *is uniform on the triangle with vertices* $(0,\varepsilon)$, $(0,1)$ *and* $(1-\varepsilon,1)$, *where* $\varepsilon = 0.1$. *The bandwidth for the computation of the MSLE was* $b_n = n^{-1/5} \approx 0.251189$.

medical statistics. The non-separated case is rather different and has its own specific difficulties. The behavior of the MLE and SMLE in this situation is discussed in [4], [6] and [10], but the theory is still rather incomplete, even for the MLE. There is a conjecture for its asymptotic distribution, put forward in [5] and [11], but this conjecture has not been proved up till now, although a simulation study, supporting the conjecture is given in [10]. The theory for the MSLE in this situation has still not been developed.

2. Characterization of the MSLE and consistency.

Let, for an estimate $\tilde h_n$ of h_0, satisfying

$$\tilde h_n(t,u)=0,\ u-t<\varepsilon, \tag{2.1}$$

for some $\varepsilon>0$, the nabla function ∇_F be defined by:

$$\nabla_F(u)=\frac{\tilde h_{n1}(u)}{F(u)}-\frac{\tilde h_{n2}(u)}{1-F(u)}+\int_{v=0}^{u}\frac{\tilde h_n(v,u)}{F(u)-F(v)}\,dv-\int_{v=u}^{M}\frac{\tilde h_n(u,v)}{F(v)-F(u)}\,dv, \tag{2.2}$$

if $0<F(u)<1$. If $F(u)=0$ or $F(u)=1$, we define $\nabla_F(u)=0$.

Then, similarly to the ordinary MLE, the MSLE can be characterized by the so-called Fenchel duality conditions.

Lemma 2.1. *Let $\tilde h_n$ satisfy (2.1), for some $\varepsilon>0$, and let a and b be defined as in Condition 1.2 (see (1.10) and (1.11)). Then the distribution function $\hat F_n$ maximizes (1.7) if and only if $\hat F_n(t)=0$, $t<a$, $\hat F_n(t)=1$, $t>b$, and $\hat F_n$ is continuous on $[0,M)$ and satisfies the conditions*

$$\int_{v=t}^{b}\nabla_{\hat F_n}(v)\,dv\le 0,\ t\in[a,b), \tag{2.3}$$

and

$$\int_a^b\nabla_{\hat F_n}(v)\hat F_n(v)\,dv=0. \tag{2.4}$$

Moreover, if $t\in[0,M)$ is a point of increase of $\hat F_n$, i.e.,

$$\begin{cases}\hat F_n(u)-\hat F_n(u')>0, & \text{for all } u,u'\in[0,M] \text{ such that } u'<t<u, \quad , \text{ if } t>0,\\ \hat F_n(u)>0, & u\in(0,M] \quad , \text{ if } t=0,\end{cases} \tag{2.5}$$

we have:

$$\nabla_{\hat F_n}(t)=0 \qquad \text{and} \qquad \int_t^b\nabla_{\hat F_n}(v)\,dv=0. \tag{2.6}$$

Proof. It is clear that if $\hat F_n$ gives mass to $[0,a)$ or $[b,M]$, we can improve $\ell(F)$ by displacing mass to the interval $[a,b)$. Therefore, we may assume that $\hat F_n(t)=0$, $t<a$, $\hat F_n(t)=1$, $t>b$.

Now first suppose that the conditions (2.3) and (2.4) are satisfied. Then we cannot have $\hat F_n(t)=0$ for t in an interval where $\tilde h_{n1}(t)>0$ or $\int_{u=0}^{t}\tilde h_n(u,t)>0$, since otherwise $\ell(\hat F_n)=-\infty$. Similarly, we cannot have $\hat F_n(t)=1$ for t in an interval where $\tilde h_{n2}(t)>0$ or $\int_{u=t}^{M}\tilde h_n(t,u)>0$.

Since the criterion function $F\mapsto\ell(F)$ is concave in F, we get:

$$\ell(F)-\ell(\hat F_n)\le\int_0^M\nabla_{\hat F_n}(u)\{F(u)-\hat F_n(u)\}\,du=\int_a^b\nabla_{\hat F_n}(u)\{F(u)-\hat F_n(u)\}\,du, \tag{2.7}$$

where we use the facts that the integrals defining $\ell(F)$ are all non-positive and that $\nabla_{\hat F_n}(t)=0$, if $t\notin[a,b)$. Note that this is similar to the relation (1.11) in [11].

By (2.4),

$$\int_a^b \nabla_{\hat F_n}(u)\hat F_n(u)\,du=0,$$

and hence:

$$\int_a^b \nabla_{\hat F_n}(u)\{F(u)-\hat F_n(u)\}\,du=\int_a^b \nabla_{\hat F_n}(u)F(u)\,du.$$

If $F=1_{[t,\infty)}$ for some $t\in[a,b)$, we get by (2.3),

$$\int_a^b \nabla_{\hat F_n}(u)F(u)\,du=\int_t^b \nabla_{\hat F_n}(u)\,du\le 0.$$

So we also get, for distribution functions of the type

$$F=\sum_{i=1}^k \alpha_i 1_{[t_i,\infty)},\ a\le t_1<\ldots,t_k\le b,\ \alpha_i\in(0,1),\ \sum_{i=1}^k\alpha_i\le 1,$$

that

$$\int_a^b \nabla_{\hat F_n}(u)F(u)\,du=\sum_{i=1}^k\alpha_i\int_{t_i}^b \nabla_{\hat F_n}(u)\,du\le 0.$$

Since we can approximate any distribution function F on $[a,b]$ by distribution functions of this type, this implies $\ell(F)\le\ell(\hat F_n)$, for all distribution functions F.

Conversely, suppose that $\hat F_n$ maximizes $\ell(F)$. Then we must have, if $t\in(a,b)$ and $F=1_{[t,\infty)}$,

$$\int_{v=t}^b \nabla_{\hat F_n}(v)\,dv=\lim_{\delta\downarrow 0}\delta^{-1}\bigl\{\ell(\hat F_n+\delta F)-\ell(\hat F_n)\bigr\}\le 0,$$

(using the concavity of ℓ for the existence of the limit), and hence (2.3) has to be satisfied for $\hat F_n$. Moreover, defining F_δ by

$$F_\delta(t)=(1+\delta)\hat F_n(t)\wedge 1,\ t\in[0,M],$$

we find:

$$\lim_{\delta\to 0}\frac{\ell(F_\delta)-\ell(\hat F_n)}{\delta}=0,$$

since the limit has to be nonpositive, if we let δ tend to zero, either from above or from below.

We have:

$$\lim_{\delta\to 0}\frac{\ell(F_\delta)-\ell(\hat F_n)}{\delta}=\int_a^b \nabla_{\hat F_n}(u)\hat F_n(u)\,du,$$

so (2.4) must hold.

Suppose $\hat F_n$ has a jump at $t\in(a,b)$ and suppose $\nabla_{\hat F_n}(t-)>0$. Define

$$F_\delta(u)=\begin{cases}\hat F_n(u), & u<t-\delta,\\ \hat F_n(t), & u\in[t-\delta,t)\\ \hat F_n(u), & u\in[t,M].\end{cases}$$

Then

$$\int \nabla_{\hat F_n}(u)\{F_\delta(u)-\hat F_n(u)\}\,du>0,$$

for small $\delta>0$, a contradiction. Hence we must have: $\nabla_{\hat F_n}(t-)\le 0$. If $\nabla_{\hat F_n}(t)<0$, we define

$$F_\delta(u)=\begin{cases}\hat F_n(u), & u<t,\\ \hat F_n(t-), & u\in[t,t+\delta)\\ \hat F_n(u), & u\in[t+\delta,M],\end{cases}$$

and then again:

$$\int \nabla_{\hat F_n}(u)\{F_\delta(u)-\hat F_n(u)\}\,du>0,$$

for small $\delta>0$, a contradiction, so we must have: $\nabla_{\hat F_n}(t)\ge 0$, implying we must have

$$\nabla_{\hat F_n}(t-)\le 0\le \nabla_{\hat F_n}(t). \tag{2.8}$$

On the other hand, we have by the continuity of $\tilde h_{nj}$, $j=1,2$, and $\tilde h_n$ and the definition of a and b:

$$\begin{aligned}\nabla_{\hat F_n}(t)-\nabla_{\hat F_n}(t-)&=\frac{\tilde h_{n1}(t)}{\hat F_n(t)}-\frac{\tilde h_{n1}(t)}{\hat F_n(t-)}-\frac{\tilde h_{n2}(t)}{1-\hat F_n(t)}+\frac{\tilde h_{n2}(t)}{1-\hat F_n(t-)}\\&\quad+\int_{v=0}^{u}\frac{\tilde h_n(v,t)}{\hat F_n(t)-\hat F_n(v)}\,dv-\int_{v=0}^{u}\frac{\tilde h_n(v,t)}{\hat F_n(t-)-\hat F_n(v)}\,dv\\&\quad-\int_{v=t}^{M}\frac{\tilde h_n(t,v)}{\hat F_n(v)-\hat F_n(t)}\,dv+\int_{v=t}^{M}\frac{\tilde h_n(t,v)}{\hat F_n(v)-\hat F_n(t-)}\,dv<0,\end{aligned}$$

contradicting (2.8). The conclusion is that we must have: $\hat F_n(t-)=\hat F_n(t)$, for $t\in(a,b)$.

If $t=a$, we either have: $a=0$, in which case $\hat F_n$ is continuous at a by right-continuity, or we have $\hat F_n(t)=0$, $t<a$. Then, if $\hat F_n$ would have a jump at a, we define the perturbation

$$F(u)=\begin{cases}0, & u<a,\\ \hat F_n(a)/2, & u\in[a,a+\delta)\\ \hat F_n(u), & u\in[a+\delta,M],\end{cases}$$

Then we get, using condition (1.13),

$$\nabla_{\hat F_n}(a)\{F_\delta(a)-\hat F_n(a)\}=\frac{\hat F_{n1}(a)\tilde h_{n2}(a)}{2\{1-\hat F_n(a)\}}+\tfrac12\hat F_n(a)\int_{v=t}^{M}\frac{\tilde h_n(t,v)}{\hat F_n(v)-\hat F_n(t)}\,dv>0$$

Hence

$$\lim_{\delta\downarrow 0}\delta^{-1}\left\{\ell(F_\delta)-\ell(\hat F_n)\right\}=\frac{\hat F_{n1}(a)\tilde h_{n2}(a)}{2\{1-\hat F_n(a)\}}+\tfrac12\hat F_n(a)\int_{v=t}^{M}\frac{\tilde h_n(t,v)}{\hat F_n(v)-\hat F_n(t)}\,dv>0,$$

contradicting the assumption that $\hat F_n$ maximizes $\ell(F)$. So we must have $\hat F_n(a)=0$. Similarly, we either have $b=M$ or we have $b<M$ and $\hat F_n(b-)=1$.

Finally, suppose (2.5) is satisfied for a point $t \in (a,b)$ and suppose $\nabla_{\hat{F}_n}(t) > 0$. Then, by the continuity of $\hat{F}_n$, there also exists a neighborhood of t such that $\nabla_{\hat{F}_n}(u) > 0$ for u in this neighborhood. We now define a perturbation F_δ of $\hat{F}_n$ by:

$$F_\delta(u) = \begin{cases} \hat{F}_n(u), & u < t-\delta, \\ \hat{F}_n(t+\delta), & u \in [t-\delta, t+\delta) \\ \hat{F}_n(u), & u \in [t+\delta, M]. \end{cases}$$

Then we have for sufficiently small $\delta > 0$:

$$\int \nabla_{\hat{F}_n}(u)\{F_\delta(u) - \hat{F}_n(u)\}\,du > 0,$$

contradicting

$$\int \nabla_{\hat{F}_n}(u)\{F_\delta(u) - \hat{F}_n(u)\}\,du \le 0.$$

If (2.5) is satisfied for a point $t \in (a,b)$ and $\nabla_{\hat{F}_n}(t) < 0$, we define the perturbation F_δ of $\hat{F}_n$ by:

$$F_\delta(u) = \begin{cases} \hat{F}_n(u), & u < t-\delta, \\ \hat{F}_n(t-\delta), & u \in [t-\delta, t+\delta) \\ \hat{F}_n(u), & u \in [t+\delta, M]. \end{cases}$$

and get a contradiction in the same way. So, if (2.5) is satisfied for a point $t \in (a,b)$, we must have:

$$\nabla_{\hat{F}_n}(t) = 0.$$

If a is a point of increase (which can only happen if $\hat{F}_n(u) > 0$ for $u > a$), we argue in the same way, taking a right neighborhood of a. Since $\hat{F}_n$ cannot have points of increase t such that $0 < t < a$ or $b < t < M$, this proves the left-hand side of (2.6).

Furthermore, $t \in [a,b)$ can only satisfy (2.5) if

$$\begin{aligned} \int_a^b \nabla_{\hat{F}_n}(u)\hat{F}_n(u)\,du &= \left[-\hat{F}_n(t)\int_{u=t}^b \nabla_{\hat{F}_n}(u)\,du\right]_{t=a}^b + \int_{t=a}^b\int_{u=t}^b \nabla_{\hat{F}_n}(u)\,du\,d\hat{F}_n(t) \\ &= \hat{F}_n(a)\int_{u=a}^b \nabla_{\hat{F}_n}(u)\,du + \int_{t=a}^b\int_{u=t}^b \nabla_{\hat{F}_n}(u)\,du\,d\hat{F}_n(t) \\ &= \int_{t=a}^b\int_{u=t}^b \nabla_{\hat{F}_n}(u)\,du\,d\hat{F}_n(t), \end{aligned} \tag{2.9}$$

using the second equality in (2.4), implying by (2.3) that

$$\int_{u=t}^b \nabla_{\hat{F}_n}(u)\,du = 0,$$

for points t satisfying (2.5), since otherwise the right-hand side of (2.9) would be strictly negative. □

Note that if $\nabla_F(t) = 0$ for all $t \in (a,b)$, where ∇_F is defined by (2.2), the conditions of Lemma 2.1 are satisfied for F, and hence F would be the MSLE if it also would be a distribution function.

But unfortunately, the function F satisfying $\nabla_F(t) = 0$ for all $t \in (a,b)$, although usually uniquely determined, need not be monotone. We will call a function $\tilde{F}_n$, satisfying $\nabla_{\tilde{F}_n}(t) = 0$, $t \in (a,b)$, a *plug-in estimator* or *naive estimator* (as in [8]). This plug-in estimator will be further studied in section 4 in the proof of the local asymptotic normality of the MSLE, where it will be shown that the MSLE is indeed locally asymptotically equivalent to this plug-in estimator.

COROLLARY 2.1. *Let $\tilde{h}_n$ satisfy (2.1), for some $\varepsilon > 0$, and let a and b be defined as in Lemma 2.1. Then the distribution function $\hat{F}_n$ maximizes (1.7) if and only if $\hat{F}_n(t) = 0$, $t < a$, $\hat{F}_n(t) = 1$, $t > b$, and $\hat{F}_n$ is continuous on $[0,M)$ and satisfies the conditions*

$$\int_a^t \nabla_{\hat{F}_n}(v)\,dv \ge 0,\ t \in (a,b), \tag{2.10}$$

and

$$\int_a^b \nabla_{\hat{F}_n}(v)\,dv = 0. \tag{2.11}$$

Moreover, if $t \in [0,M)$ is a point of increase of $\hat{F}_n$, i.e., satisfies condition (2.5) of Lemma 2.1, then

$$\nabla_{\hat{F}_n}(t) = 0 \qquad \textit{and} \qquad \int_a^t \nabla_{\hat{F}_n}(v)\,dv = 0. \tag{2.12}$$

Proof. Suppose $\hat{F}_n$ maximizes $\ell(F)$. Defining

$$F_\delta(t) = \{1 - (1+\delta)(1 - F(t))\} \vee 0,\ t \in [0,M],$$

we find:

$$\lim_{\delta \to 0} \frac{\ell(F_\delta) - \ell(\hat{F}_n)}{\delta} = \int_a^b \nabla_{\hat{F}_n}(u)\big\{1 - \hat{F}_n(u)\big\}\,du = 0. \tag{2.13}$$

So if $\hat{F}_n$ maximizes $\ell(F)$, (2.11) follows from (2.13) and (2.4) of Lemma 2.1.

$$\int_a^b \nabla_{\hat{F}_n}(u)\,du = 0.$$

This implies:

$$\int_a^t \nabla_{\hat{F}_n}(v)\,dv = -\int_t^b \nabla_{\hat{F}_n}(v)\,dv,$$

and conditions (2.4) and (2.6) now also follow. Conversely, if the conditions of the corollary hold, we get:

$$\begin{aligned}\int_a^b \hat{F}_n(u)\nabla_{\hat{F}_n}(u)\,du &= \hat{F}_n(b)\int_a^b \nabla_{\hat{F}_n}(v)\,dv + \int_{t=a}^b \int_{v=a}^u \nabla_{\hat{F}_n}(v)\,dv\,d\hat{F}_n(u)\\ &= \hat{F}_n(b)\int_a^b \nabla_{\hat{F}_n}(v)\,dv = 0,\end{aligned}$$

implying condition (2.4) of Lemma 2.1. The other conditions of 2.1 follow similarly. □

We now simplify the conditions somewhat, in view of the iterative convex minorant agorithm, to be discussed in section 3. Multiplying ∇_F by $F(1-F)$ yields the function

$$\begin{aligned}\overline{\nabla}_F(u) &= \tilde h_{n1}(u)\{1-F(u)\} - \tilde h_{n2}(u)F(u)\\ &\qquad + F(u)\{1-F(u)\}\left\{\int_{v=0}^{u}\frac{\tilde h_n(v,u)}{F(u)-F(v)}\,dv - \int_{v=u}^{M}\frac{\tilde h_n(u,v)}{F(v)-F(u)}\,dv\right\}. \end{aligned}\tag{2.14}$$

Corollary 2.2. *Let $\tilde h_n$ satisfy (2.1), for some $\varepsilon > 0$ and let the function $\overline{\nabla}_F$ be defined by (2.14). Then the distribution function $\hat F_n$ maximizes (1.7) if and only if $\hat F_n(t) = 0$, $t < a$, $\hat F_n(t) = 1$, $t > b$, and $\hat F_n$ is continuous on $[0, M)$ and satisfies the conditions*

$$\int_0^t \overline{\nabla}_{\hat F_n}(v)\,dv \ge 0,\ t \in [0,M],\tag{2.15}$$

and

$$\int_0^M \overline{\nabla}_{\hat F_n}(v)\,dv = 0.\tag{2.16}$$

Moreover, if $t \in [0, M)$ is a point of increase of $\hat F_n$, i.e., satisfies condition (2.5) of Lemma 2.1, then

$$\overline{\nabla}_{\hat F_n}(t) = 0 \qquad \text{and} \qquad \int_0^t \overline{\nabla}_{\hat F_n}(v)\,dv = 0.\tag{2.17}$$

Proof. Let a and b be defined as in Corollary 2.1. We then have, for $t \in [a, b)$,

$$\int_0^t \overline{\nabla}_{\hat F_n}(v)\,dv = \int_a^t \hat F_n(u)\{1-\hat F_n(u)\}\nabla_{\hat F_n}(v)\,dv$$

We have:

$$\begin{aligned}&\int_a^t \hat F_n(u)\{1-\hat F_n(u)\}\nabla_{\hat F_n}(v)\,dv\\ &= \hat F_n(t)\{1-\hat F_n(t)\}\int_{u=a}^t \nabla_{\hat F_n}(u)\,du - \int_a^t\{1-2\hat F_n(u)\}\int_{v=a}^u \nabla_{\hat F_n}(v)\,dv\,d\hat F_n(u)\\ &= \hat F_n(t)\{1-\hat F_n(t)\}\int_{u=a}^t \nabla_{\hat F_n}(u)\,du.\end{aligned}$$

Hence condition (2.15) is equivalent to condition (2.10) of Corollary 2.1. Similarly, by Corollary 2.1, condition (2.15) can be written

$$\int_0^M \overline{\nabla}_{\hat F_n}(v)\,dv = \int_a^b \hat F_n(u)\{1-\hat F_n(u)\}\nabla_{\hat F_n}(v)\,dv = \hat F_n(b)\{1-\hat F_n(b)\}\int_{u=a}^b \nabla_{\hat F_n}(u)\,du,$$

and is therefore equivalent to condition (2.10) of Corollary 2.1. □

We now prove consistency of the MSLE. The proof is somewhat analogous to the proof of the consistency of the MLE in [11].

THEOREM 2.1. **(Consistency of the MSLE)** *Let Condition 1.1 be satisfied on* $[0, M]$ *for the distribution function* F_0 *and the observation density* g*. Moreover, let* $\tilde h_{nj}$ *and* $\tilde h_n$ *be kernel estimators of* h_{0j} *and* h_0*, respectively, of the type defined in Condition 1.1. Finally, let* $\hat F_n$ *be the MSLE of* F_0*. Then, with probability one,*

$$\lim_{n\to\infty} \hat F_n(t) = F_0(t),$$

for each $t \in [0, M)$*. The convergence is uniform on each subinterval* $[a, b]$ *of* $(0, M)$*.*

Proof. By the assumption on the kernel estimates and the observation density g, we may assume that $\tilde h_n$ satisfies (2.1), for some $\varepsilon > 0$, and all large n. Let the function ψ be defined by

$$\psi(F; h_1, h_2, h) = \int h_1(t)\log F(t)\,dt + \int h_2(t)\log\{1 - F(t)\}\,dt + \int h(t,u)\log\{F(u) - F(t)\}\,dt\,du, \tag{2.18}$$

Then we must have, if $h_j = \tilde h_{nj}$, $j = 1, 2$, and $h = \tilde h_n$,

$$\lim_{\varepsilon\downarrow 0} \varepsilon^{-1}\left\{\psi\big((1-\varepsilon)\hat F_n + \varepsilon F_0; h_1, h_2, h\big) - \psi(\hat F_n; h_1, h_2, h)\right\} \le 0.$$

This implies (see also (4.20) in [11]):

$$\int \frac{F_0(t)}{\hat F_n(t)}\tilde h_{n1}(t)\,dt + \int \frac{1 - F_0(t)}{1 - \hat F_n(t)}\tilde h_{n2}(t)\,dt + \int \frac{F_0(u) - F_0(t)}{\hat F_n(u) - \hat F_n(t)}\tilde h_n(t,u)\,dt\,du \le 1. \tag{2.19}$$

Fix a small $\delta \in [0, M/2]$ and let the intervals A_δ and B_δ be defined by

$$A_\delta = [\delta, M], \qquad B_\delta = [0, M - \delta].$$

Moreover, if $\{(t,u) : g(t,u) > 0\} = \{(t,u) : u - t > \varepsilon\}$, let C_δ be defined by

$$C_\delta = \{(t,u) : t < u,\ d((t,u),(t,\gamma(t)) \ge \delta\},$$

where γ is the function defining the boundary of the support of g, as in Condition 1.1, and d denotes the Euclidean distance. Then, arguing as in [11], Part II, Chapter 4, we find that there exists an $M > 0$ such that for all n,

$$\sup_{t\in A_\delta} 1/\hat F_n(t) + \sup_{t\in B_\delta} 1/\{1 - \hat F_n(t)\} \le M.$$

We also cannot have that $\hat F_n(u_{k_n}) - \hat F_n(t_{k_n}) \to 0$, for a sequence of points $(t_{k_n}, u_{k_n}) \in C_\delta$. For suppose, if necessary by taking a subsequence, that $t_{k_n} \to t_0$ and $u_{k_n} \to u_0$. The $u_0 - t_0 \ge \varepsilon + \delta$. By the vague convergence of $\hat F_n$ to F, there are continuity points t_1 and u_1 such that $t_0 < t_1 < u_1 < u_0$, $u_1 - t_1 \ge \frac{1}{2}\delta + \varepsilon$, and $\hat F_n(t_1) \to F(t_1)$ and $\hat F_n(u_1) \to F(u_1)$. Moreover, since

$$\hat F_n(u_1) - \hat F_n(t_1) \le \hat F_n(u_{k_n}) - \hat F_n(t_{k_n}),$$

for large n, we must have: $F(u_1) - F(t_1) = 0$. We then would get that there exists a rectangle $[t_1, t_2] \times [u_2, u_1]$ such that $u_2 - t_2 > \varepsilon$ and

$$\begin{aligned}\liminf_{n\to\infty} &\int_{[t_1,t_2]\times[u_2,u_1]} \frac{F_0(u) - F_0(t)}{\hat F_n(u) - \hat F_n(t)}\tilde h_n(t,u)\,dt\,du\\ &\ge K\{F_0(u_2) - F_0(t_2)\}\int_{[t_1,t_2]\times[u_2,u_1]} h_0(t,u)\,dt\,du,\end{aligned}$$

for any $K > 0$, contradicting (2.19). So we may also assume that

$$\inf_{(t,u)\in C_\delta}\{\hat F_n(u)-\hat F_n(t)\}\ge \frac{1}{M}\,,$$

for all n.

As in [11], Part II, Chapter 4, we have by the Helly compactness theorem that the exists a set of probability one, such that for each ω in this set the sequence $(\hat F_n(\cdot;\omega))$ has a subsequence $\hat F_{n_k}(\cdot;\omega)$ which converges vaguely to a subdistribution function $F = F(\cdot;\omega)$. By the vague convergence of $\hat F_{n_k}(\cdot;\omega)$ to F we now get:

$$\begin{aligned}&\int_{A_\delta}\frac{F_0(t)}{\hat F_{n_k}(t;\omega)}\tilde h_{n1}(t)\,dt+\int_{B_\delta}\frac{1-F_0(t)}{1-\hat F_{n_k}(t;\omega)}\tilde h_{n2}(t)\,dt+\int_{C_\delta}\frac{F_0(u)-F_0(t)}{\hat F_{n_k}(u;\omega)-\hat F_{n_k}(t;\omega)}\tilde h_n(t,u)\,dt\,du\\&\to\int_{A_\delta}\frac{F_0(t)}{F(t)}h_{01}(t)\,dt+\int_{B_\delta}\frac{1-F_0(t)}{1-F(t)}h_{02}(t)\,dt+\int_{C_\delta}\frac{F_0(u)-F_0(t)}{F(u)-F(t)}h_0(t,u)\,dt\,du,\ n\to\infty.\end{aligned}$$

By monotone convergence we now also have:

$$\begin{aligned}&\int_{[0,M]}\frac{F_0(t)}{F(t)}h_{01}(t)\,dt+\int_{[0,M]}\frac{1-F_0(t)}{1-F(t)}h_{02}(t)\,dt+\int_{[0,M]^2}\frac{F_0(u)-F_0(t)}{F(u)-F(t)}h_0(t,u)\,dt\,du\\&=\lim_{\delta\downarrow0}\left\{\int_{A_\delta}\frac{F_0(t)}{F(t)}h_{01}(t)\,dt+\int_{B_\delta}\frac{1-F_0(t)}{1-F(t)}h_{02}(t)\,dt+\int_{C_\delta}\frac{F_0(u)-F_0(t)}{F(u)-F(t)}h_0(t,u)\,dt\,du\right\}\le 1.\end{aligned}\tag{2.20}$$

Suppose $F(t)\ne F_0(t)$ for some $t\in[0,M/2]$. Then there exist a $u\in(t,M)$ such that $h_0(t,u)>0$ and

$$\frac{F_0(t)^2}{F(t)}+\frac{\{1-F_0(u)\}^2}{1-F(u)}+\frac{\{F_0(u)-F_0(t)\}^2}{F(u)-F(t)}>1,$$

since

$$\frac{F_0(t)^2}{x}+\frac{\{1-F_0(u)\}^2}{1-y}+\frac{\{F_0(u)-F_0(t)\}^2}{y-x}\begin{cases}=1, & F_0(t)=x,\ F_0(u)=y,\\ >1, & \text{otherwise.}\end{cases}$$

(see also (4.27) in [11]). By the continuity of F_0 and the monotonicity and right continuity of F there exist therefore also $h>0$ such that

$$\frac{F_0(t')^2}{F(t')}+\frac{\{1-F_0(u')\}^2}{1-F(u')}+\frac{\{F_0(u')-F_0(t')\}^2}{F(u')-F(t')}>1,$$

if $t'\in[t,t+h]$ and $u'\in[u,u+h]$. This implies

$$\begin{aligned}&\int_{[0,M]}\frac{F_0(t)}{F(t)}h_{01}(t)\,dt+\int_{[0,M]}\frac{1-F_0(t)}{1-F(t)}h_{02}(t)\,dt+\int_{[0,M]^2}\frac{F_0(u)-F_0(t)}{F(u)-F(t)}h_0(t,u)\,dt\,du\\&=\int_{[0,M]}\frac{F_0(t)^2}{F(t)}g_1(t)\,dt+\int_{[0,M]}\frac{\{1-F_0(t)\}^2}{1-F(t)}g_2(t)\,dt+\int_{[0,M]^2}\frac{\{F_0(u)-F_0(t)\}^2}{F(u)-F(t)}g(t,u)\,dt\,du\\&=\int_{[0,M]^2}\left\{\frac{F_0(t)^2}{F(t)}+\frac{\{1-F_0(t)\}^2}{1-F(t)}+\frac{\{F_0(u)-F_0(t)\}^2}{F(u)-F(t)}\right\}g(t,u)\,dt\,du>1,\end{aligned}$$

in contradiction with (2.20). So we must have $F(t)=F_0(t)$ if $t\in[0,M/2]$. A similar argument yields $F(t)=F_0(t)$ if $t\in[M/2,M)$.

So, for each ω outside a set of probability zero, the sequence $(\hat F_n(\cdot;\omega))$ has a subsequence which converges weakly to F_0. This implies that $\hat F_n(t)$ converges almost surely to $F_0(t)$ for each $t\in[0,M)$. The uniformity of the convergence on subintervals follows from the continuity of F_0. □

3. Algorithms. We start by recalling how the MSLE is computed for current status data. As explained in [8], the MSLE can be computed by first constructing the continuous cusum diagram

$$\left(\tilde G_n(t),\tilde H_n(t)\right),\, t\in[0,M], \tag{3.1}$$

defining the densities $\tilde g_n$ and $\tilde h_n$ on $[0,M]$ by

$$\tilde g_n(t)=\frac1n\sum_{i=1}^n K_{b_N}(t-T_i),\qquad \tilde h_n(t)=\frac1n\sum_{i=1}^n K_{b_N}(t-T_i)\Delta_i, \tag{3.2}$$

and $\tilde G_n$ and $\tilde H_n$ by:

$$\tilde G_n(t)=\int_0^t \tilde g_n(u)\,du,\qquad \tilde H_n(t)=\int_0^t \tilde h_n(u)\,du,\, t\in[0,M],$$

where

$$\Delta_i=1_{\{X_i\le T_i\}},$$

noting that for the current status model we just have the observation times T_i and the information whether the unobervable event of interest X_i is to the left or to the right of T_i.

Then the slope of the greatest convex minorant of the cusum diagram (3.1) defines the MSLE $\hat F_n$ and $\hat F_n$ maximizes the smoothed log likelihood

$$\int_0^M \tilde h_n(t)\log F(t)\,dt+\int_0^M\left\{\tilde g_n(t)-\tilde h_n(t)\right\}\log\{1-F(t)\}\,dt.$$

One can improve the estimates (3.2) by using boundary kernels, but this is not relevant here, the main point is that the MSLE can in this case be computed by a one-step algorithm which just computes the greatest convex minorant of the cusum diagram (3.1). This parallels the situation for the MLE in the current status model, which can also be computed by constructing a cusum diagram and the corresponding convex minorant. Such a simple one-step algorithm does no longer exist for the computation of the MLE for the interval censoring, case 2, model, and the situation is similar for the MSLE: we have to use some iterative procedure.

The EM algorithm is based on the following “self-consistency equations”

$$\hat f_n(t)=\left\{\int_t^M\frac{\tilde h_{n1}(v)}{\hat F_n(v)}\,dv+\int_0^t\frac{\tilde h_{n2}(v)}{1-\hat F_n(v)}\,dv+\int_{v<t<u}\frac{\tilde h_n(v,u)}{\hat F_n(u)-\hat F_n(v)}\,dv\,du\right\}\hat f_n(t),$$

where $\hat f_n(t)=\hat F_n'(t)$. This yields the iteration steps

$$f^{(k+1)}(t)=\left\{\int_t^M\frac{\tilde h_{n1}(v)}{F^{(k)}(v)}\,dv+\int_0^t\frac{\tilde h_{n2}(v)}{1-F^{(k)}(v)}\,dv+\int_{v<t<u}\frac{\tilde h_n(v,u)}{F^{(k)}(u)-F^{(k)}(v)}\,dv\,du\right\}f^{(k)}(t). \tag{3.3}$$

One can indeed use a discretized version of (3.3) to compute the MSLE, but the EM algorithm is (as is usual for this type of problem with many parameters) very slow. Simply enhancing the EM algorithm by a Newton step is also not helpful because of the many constraints the solution has to satisfy, leading to very small “feasible steps”. For this reason a Newton-improved EM algorithm does not improve very much on the EM algorithm itself.

An algorithm that we found very stable and rather fast is the interior point method with a logarithmic barrier function, which keeps the path to the solution away from the boundary. In this case one maximizes the criterion function

$$\begin{aligned}\phi_\mu(F) &= \sum_{i=1}^m \{\tilde h_{n1}(t_i)\log F(t_i)\}d_i + \sum_{i=1}^m \{\tilde h_{n2}(t_i)\log\{1-F(t_i)\}\}d_i\\ &+ \sum_{i=1}^{m-1}\sum_{j=i+1}^m \{\tilde h_n(t_i,t_j)\log\{F(t_j)-F(t_i)\}\}d_id_j + \mu\sum_{i=1}^{m+1}\log\{F(t_i)-F(t_{i-1})\},\end{aligned} \tag{3.4}$$

instead of (1.8), where $F(t_0)=0$ and $F(t_{m+1})=1$, and lets the parameter μ tend to zero from above after completing an interior iteration loop, maximizing (3.4) for a fixed μ. In the interior iteration loop one uses the Hessian matrix, corresponding to (3.4), and solves the maximization problem by doing Newton-Raphson steps in maximizing (3.4). The iterations are started with $\mu = 1$ and μ is halved after completing an interior iteration after which (3.4) is maximized again, starting with the value of F, obtained in the preceding inner iteration.

In our experience the fastest algorithm, however, is a combination of the EM algorithm with a version of the iterative convex minorant (ICM) algorithm, introduced in [5] and [11]. This procedure has the added advantage that one does not have to solve a rather large matrix equation, using the Hessian matrix, as is needed in the interior point method. Instead of the cusum diagram (3.1) we use a sequence of cusum diagrams

$$\left(W_n^{(k)}(t), V_n^{(k)}(t)\right),\ t\in[0,M],\ k=0,1,2,\dots, \tag{3.5}$$

for which we compute the greatest convex minorants at each kth step. We alternate this with an EM-step (the combination is sometimes called the "hybrid algorithm"). The cumulative weight function $W_n^{(k)}$ is of the form

$$W_n^{(k)} = \int_0^t w_n^{(k)}(u)\,du,\ t\ge 0,$$

for suitably (but somewhat arbitrarily) chosen weights $w_n^{(k)}$, and the cusum function $V_n^{(k)}$ is of the form:

$$V_n^{(k)}(t) = \int_0^t F^{(k)}(u)w_n^{(k)}(u)\,du + \int_0^t \overline{\nabla}_{F^{(k)}}(u)\,du,\ t\ge 0,$$

where, for a distribution function F, $\overline{\nabla}_F$ is the function, defined by (2.14), evaluated at $F=F^{(k)}$. The idea is that the iterations force the Fenchel duality conditions (2.15) and (2.16) to be satisfied at the end of the iterations.

The following weight function, chosen by taking the diagonal elements of the Hessian matrix, corresponding to the function $\overline{\nabla}_F$, gave good convergence results in our simulation study of the MSLE:

$$\begin{aligned}w_n^{(k)}(t) &= \tilde h_{n1}(t) + \tilde h_{n2}(t)\\ &\quad - \{1-2F_n^{(k)}(t)\}\left\{\int_{u=0}^t \frac{\tilde h_n(u,t)}{F^{(k)}(t)-F^{(k)}(u)}\,du - \int_{u=t}^M \frac{\tilde h_n(t,u)}{F^{(k)}(u)-F^{(k)}(t)}\,du\right\}\\ &\quad + \int_{u=0}^t \frac{\tilde h_n(u,t)}{\{F^{(k)}(t)-F^{(k)}(u)\}^2}\,du + \int_{u=t}^M \frac{\tilde h_n(t,u)}{\{F^{(k)}(u)-F^{(k)}(t)\}^2}\,du.\end{aligned}$$

To prevent divergence of the algorithm, Armijo's line search method, as implemented in [12], was used for determining the step size at each iteration.

Note that, in the case of current status data, the function $\overline{\nabla}_F$ is just given by:

$$\overline{\nabla}_F = \tilde{h}_n(t) - \tilde{g}_n(t)F(t)\,, \tag{3.6}$$

and that we can compute $\hat{F}_n$ in one step this case, using the cusum diagram (3.1).

4. Asymptotic distribution.

In this section we will prove the following theorem.

THEOREM 4.1. *Let condition 1.1 be satisfied. Moreover, let F_0 be twice differentiable, with a bounded continuous derivative f_0 on the interior of $[0,M]$, which is bounded away from zero on $[0,M]$, with a finite positive right limit at 0 and a positive left limit at M. Also, let f_0 have a bounded continuous derivative on $(0,M)$ and let g_1 and g_2 be twice differentiable on the interior of their supports S_1 and S_2, respectively, and let $g_1F_0+g_2(1-F_0)$ stay away from zero on $[0,M]$. Furthermore, let the joint density g of the pair of observation times (T_i,U_i) have a bounded (total) second derivative on the interior of its support S, having finite limits approaching the boundary of S. Suppose that X_i is independent of (T_i,U_i), and let d_{F_0} be defined by*

$$d_{F_0}(v) = \frac{F_0(v)\{1-F_0(v)\}}{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)}\,.$$

Then, if $b_n \asymp n^{-1/5}$, we have for for each $v\in(0,M)$,

$$\sqrt{nb_n}\left\{\hat{F}_n(v)-F_0(v)-\frac{\beta(v)b_n^2}{2\sigma_1(v)}\right\} \xrightarrow{\mathcal{D}} N\left(0,\sigma(v)^2\right),$$

where

$$\begin{aligned}\beta(v) &= \frac{\{1-F_0(v)\}h_1''(v)-F_0(v)h_2''(v)}{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)}\int u^2K(u)\,du\\ &\quad + d_{F_0}(v)\left\{\int_{t=0}^{v}\frac{\frac{\partial^2}{\partial v^2}h_0(t,v)}{F_0(v)-F_0(t)}\,dt-\int_{u=v}^{M}\frac{\frac{\partial^2}{\partial v^2}h_0(v,u)}{F_0(u)-F_0(v)}\,du\right\}\int u^2K(u)\,du,\end{aligned} \tag{4.1}$$

and

$$\sigma_1(v) = 1+d_{F_0}(v)\left\{\int_{t<v}\frac{g(t,v)}{F_0(v)-F_0(t)}\,dt+\int_{w>v}\frac{g(v,w)}{F_0(w)-F_0(v)}\,dw\right\}, \tag{4.2}$$

and where $N\left(0,\sigma(v)^2\right)$ is a normal distribution with first moment zero and variance $\sigma(v)^2$, defined by

$$\sigma(v)^2 = \frac{d_{F_0}(v)}{\sigma_1(v)}\int K(u)^2\,du. \tag{4.3}$$

Before embarking on the proof, we motivate the approach taken. Our starting point is given by the duality conditions (2.15) and (2.16). It is clear that if we would have equality in (2.15) instead of inequality, we would get the following relation by differentiating w.r.t. t:

$$\begin{aligned}\overline{\nabla}_F(t) &= \tilde{h}_{n1}(t)\{1-F(t)\}-\tilde{h}_{n2}(t)F(t)\\ &\quad + F(t)\{1-F(t)\}\left\{\int_{v=0}^{t}\frac{\tilde{h}_n(v,t)}{F(t)-F(v)}\,dv-\int_{v=t}^{M}\frac{\tilde{h}_n(t,v)}{F(v)-F(t)}\,dv\right\}=0.\end{aligned} \tag{4.4}$$

Conversely, if F solves (4.4) for each $t \in (0, M)$ and F is a distribution function, F also satisfies (2.15) and (2.16) and is therefore the MSLE.

Replacing $\tilde h_{nj}$ by h_{0j}, $j = 1, 2$, and $\tilde h_n$ by h_0, we obtain the equation

$$h_{01}(t)\{1-F(t)\} - h_{02}(t)F(t) + F(t)\{1-F(t)\}\left\{\int_{v=0}^{t}\frac{h_0(v,t)}{F(t)-F(v)}\,dv - \int_{u=t}^{M}\frac{h_0(t,u)}{F(u)-F(t)}\,du\right\} = 0,$$

which, using the definition of h_{0j} and h_0, turns into

$$\begin{aligned}&g_1(t)F_0(t)\{1-F(t)\} - g_2(t)\{1-F_0(t)\}F(t)\\ &+ F(t)\{1-F(t)\}\left\{\int_{v=0}^{t}\frac{g(v,t)\{F_0(t)-F_0(u)\}}{F(t)-F(v)}\,dv - \int_{u=t}^{M}\frac{g(t,u)\{F_0(u)-F_0(t)}{F(u)-F(t)}\,du\right\} = 0.\end{aligned}$$

This equation is clearly solved by F_0 itself.

This motivates us to consider the equation

$$\phi(t; h_1, h_2, h, F) = 0,\ t \in [0, M],$$

where

$$\begin{aligned}\phi(t; h_1, h_2, h, F) &= h_1(t)\{1-F(t)\} - h_2(t)F(t)\\ &+ F(t)\{1-F(t)\}\left\{\int_{v=0}^{t}\frac{h(v,t)}{F(t)-F(v)}\,dv - \int_{u=t}^{M}\frac{h(t,u)}{F(u)-F(t)}\,du\right\}.\end{aligned}\tag{4.5}$$

The functions h belong to a closed subset of the Banach space $C(S)$, where S is the support of g, see Condition 1.1. By Condition 1.1, S is contained in the set

$$\{(t,u) : u - t \ge \varepsilon\}$$

for a fixed $\varepsilon > 0$.

We now define the space E by

$$E = C[S_1] \times C[S_2] \times C(S) \times C[0, M],\tag{4.6}$$

where S_j if the support of g_j (see Condition 1.1), and put the following norm on E:

$$\|(h_1, h_2, h, F)\| = \max\left\{\|h_1\|, \|h_2\|, \|h\|, \|F\|\right\},\tag{4.7}$$

where the norms on the right-hand side denote the supremum norm, which we also denote by $\|\cdot\|$. Note that E is a Banach space for the norm (4.7).

We will also need another norm on $C[S]$, defined by

$$\|h\|_S = \sup_{t\in[0,M]}\left\{\int_{u:(u,t)\in S}|h(u,t)|\,du + \int_{u:(t,u)\in S}|h(t,u)|\,du\right\}.\tag{4.8}$$

Note that this is indeed a norm on $C[S]$, since $\|h\|_S = 0$ implies $h = 0$ and since the triangle inequality and homogeneity property for scalars are obviously satisfied.

LEMMA 4.1. *Let F_0, h_{01}, h_{02} and h_0 satisfy the conditions of Theorem 4.1. Furthermore, let the function ϕ be defined by (4.5). Then there exists for all small $\eta > 0$ an open set U in the Banach space $C[S_1] \times C[S_2] \times C(S)$, endowed with the norm*

$$\|(h_1, h_2, h)\| = \max\left\{\|h_1\|, \|h_2\|, \|h\|\right\},$$

such that, if $(h_1,h_2,h)\in U$, *the equation*

$$\phi(t;h_1,h_2,h,F)=0,\ t\in[0,M],$$

where ϕ *is defined by (4.5), has a unique solution* F *in the open ball* $B(F_0,\eta)\subset C[0,M]$ *with midpoint* F_0.

Proof. To avoid messy notation, we extend the definition of h to $[0,M]^2$, by defining $h(t,u)=0$, if $(t,u)\notin S$ and $h_j(t)=0$, if $t\notin S_j$. We will now use the line of argument of the proof of the implicit function theorem 10.2.1 in [1]. We define the function $\bar\phi$ by:

$$\left[\bar\phi(h_1,h_2,h,F)\right](t)=\phi(t,h_1,h_2,h,F),\ t\in[0,M], \tag{4.9}$$

so $\bar\phi$ maps E to $C[0,M]$. The derivative of $\bar\phi$ w.r.t. F, is given by the function

$$\begin{aligned}&\left[\left[\partial_4\bar\phi(h_1,h_2,h,F)\right](A)\right](t)\\&\stackrel{\text{def}}{=}-\left\{h_1(t)+h_2(t)\right\}A(t)+\{1-2F(t)\}\left\{\int_{v=0}^t\frac{h(v,t)}{F(t)-F(v)}\,dv-\int_{u=t}^M\frac{h(t,u)}{F(u)-F(t)}\,du\right\}A(t)\\&\qquad-F(t)\{1-F(t)\}\left\{\int_{u=0}^t\frac{h(u,t)\{A(t)-A(u)\}}{\{F(t)-F(u)\}^2}\,du+\int_{u=t}^M\frac{h(t,u)\{A(t)-A(u)\}}{\{F(u)-F(t)\}^2}\,du\right\},\end{aligned}$$

where $A\in C[0,M]$. Note that the right-hand side is well-defined for $F\in B(F_0,\eta)$ and small $\eta>0$, since $h(t,u)=0$ if $u-t<\varepsilon$, and since F_0 has a non-vanishing derivative on $[0,M]$, implying that $F_0(u)-F_0(t)$ stays away from zero if $u-t\ge\varepsilon$.

We now define the open set $U=U_\delta$ of functions (h_1,h_2,h) by

$$U_\delta=\left\{(h_1,h_2,h)\in C[S_1]\times C[S_2]\times C[S]:\max\left\{\|h_1-h_{01}\|,\|h_2-h_{02}\|,\|h-h_0\|_S\right\}<\delta\right\}. \tag{4.10}$$

There exists a $\delta>0$ such that for $(h_1,h_2,h)\in U_\delta$,

$$\left\|\bar\phi(h_1,h_2,h,F_1)-\bar\phi(h_1,h_2,h,F_2)-\left[\partial_4\bar\phi(h_{01},h_{02},h_0,F_0)\right](F_1-F_2)\right\|\le\varepsilon'\left\|F_1-F_2\right\|,$$

if $F_1,F_2\in B(F_0,\eta)$, where $\varepsilon'>0$ can be made arbitrarily small by making δ small.

The equation

$$\left[\partial_4\bar\phi(F_0;h_{01},h_{02},h_0,F_0)\right](A)=0$$

only has the trivial solution $A\equiv0$ in $C[0,M]$. This is seen in the following way. Suppose there exists a solution in $A\in C[0,M]$ such that $\left[\partial_4\bar\phi(F_0;h_{01},h_{02},h_0,F_0)\right](A)=0$ and $A(s)>0$ for some $s\in[0,M]$. Then also $\max_{s\in[0,M]}A(s)>0$. Suppose the maximum is attained at $t\in[0,M]$. Then:

$$\begin{aligned}&\left[\left[\partial_4\bar\phi(h_{01},h_{02},h_0,F_0)\right](A)\right](t)\\&=-\left\{h_{01}(t)+h_{02}(t)\right\}A(t)+\{1-2F_0(t)\}\left\{\int_{v=0}^t\frac{h_0(v,t)}{F_0(t)-F_0(v)}\,dv-\int_{u=t}^M\frac{h_0(t,u)}{F_0(u)-F_0(t)}\,du\right\}A(t)\\&\qquad-F_0(t)\{1-F_0(t)\}\left\{\int_{u=0}^t\frac{h_0(u,t)\{A(t)-A(u)\}}{\{F_0(t)-F_0(u)\}^2}\,du+\int_{u=t}^M\frac{h_0(t,u)\{A(t)-A(u)\}}{\{F_0(u)-F_0(t)\}^2}\,du\right\}\\&=-\left\{g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t)\right\}A(t)\\&\qquad-F_0(t)\{1-F_0(t)\}\left\{\int_{u=0}^t\frac{h_0(u,t)\{A(t)-A(u)\}}{\{F_0(t)-F_0(u)\}^2}\,du+\int_{u=t}^M\frac{h_0(t,u)\{A(t)-A(u)\}}{\{F_0(u)-F_0(t)\}^2}\,du\right\}\\&\le-\left\{g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t)\right\}A(t)<0,\end{aligned}$$

using $g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t)>0$, in contradiction with the assumption

$$[\partial_4\bar\phi(h_{01},h_{02},h_0,F_0)](A)=0.$$

We similarly get a contradiction if we assume that $A(t)<0$ for some $t\in[0,M]$ (similar arguments were used for the integral equation, studied in [2]). This shows that $\partial_4\bar\phi(h_{01},h_{02},h_0,F)$ is a linear homeomorphism of $C[0,M]$ onto $C[0,M]$ and that we can in fact use arguments of the type used in the proof of the implicit function theorem in Banach spaces, as given, for example in [1], Theorem 10.2.1.

Denoting (as in the proof of Theorem 10.2.1 of [1]) the linear mapping $\partial_4\bar\phi(h_{01},h_{02},h_0,F_0)$ by T_0 and its inverse by T_0^{-1}, we find that

$$\|T_0^{-1}\cdot\bigl\{\bar\phi(h_1,h_2,h,F_1)-\bar\phi(h_1,h_2,h,F_2)\bigr\}-\bigl(F_1-F_2\bigr)\|\le\varepsilon'\|T_0^{-1}\|\|F_1-F_2\|\le\tfrac12\|F_1-F_2\|,$$

so we have a *contraction*, and this implies that the equation

$$F=F-T_0^{-1}\cdot\bar\phi(\cdot,F;h_1,h_2,h)$$

has a unique solution $F\in B(F_0,\eta)$ which can be obtained by successive approximations, if we take the balls around h_{0j} and h_0, to which h_j and h belong, respectively, sufficiently small, using a result like 10.1.1 in [1]. This, in turn, implies that the equation

$$\bar\phi(h_1,h_2,h,F)=0$$

has a unique solution in $F\in B(F_0,\eta)$, for $(h_1,h_2,h)\in U_\delta$ and small δ. □

Having established the existence of a solution, we now consider the derivative of the solution.

LEMMA 4.2. *Let, under the conditions of Lemma 4.1, for a small $\eta>0$, $F\in B(F_0,\eta)$ be the solution of*

$$\phi(t;h_1,h_2,h,F)=0,\ t\in[0,M],$$

where ϕ is defined by (4.5), and where h_j has a bounded continuous derivative on the interior of S_j, having finite limits approaching the boundary of S_j, for $j=1,2$. Similarly, we suppose that h is differentiable on the interior of its support S and has finite limits approaching the boundary of S. Then, if $(h_1,h_2,h)\in U_\delta$, where U_δ is defined by (4.10), the solution F has a continuous and bounded derivative for sufficiently small η and δ.

Proof. If $\phi(\cdot;h_1,h_2,h,F)=0$ we have:

$$\begin{aligned}&\{1-F(t)\}h_1(t)1_{S_1}(t)-F(t)h_2(t)1_{S_2}(t)\\&\qquad+F(t)\{1-F(t)\}\left\{\int_{u:(u,t)\in S}\frac{h(u,t)}{F(t)-F(u)}\,du-\int_{u:(t,u)\in S}\frac{h(t,u)}{F(u)-F(t)}\,du\right\}=0.\end{aligned}$$

Differentiation w.r.t. t, and defining $f = F'$, yields:

$$\begin{aligned}
&\{1-F(t)\}h_1'(t)1_{S_1}(t) - F(t)h_2'(t) - f(t)\{h_1(t)1_{S_1}(t) + h_2(t)1_{S_2}(t)\}\\
&\qquad + \{1-2F(t)\}f(t)\left\{\int_{u:(u,t)\in S}\frac{h(u,t)}{F(t)-F(u)}\,du - \int_{u:(t,u)\in S}\frac{h(t,u)}{F(u)-F(t)}\,du\right\}\\
&\qquad + F(t)\{1-F(t)\}\left\{\int_{u:(u,t)\in S}\frac{\partial_2 h(u,t)}{F(t)-F(u)}\,du - \int_{u:(t,u)\in S}\frac{\partial_1 h(t,u)}{F(u)-F(t)}\,du\right\}\\
&\qquad - f(t)F(t)\{1-F(t)\}\left\{\int_{u:(u,t)\in S}\frac{h(u,t)}{\{F(t)-F(u)\}^2}\,du + \int_{u:(t,u)\in S}\frac{h(t,u)}{\{F(u)-F(t)\}^2}\,du\right\} = 0.
\end{aligned}\tag{4.11}$$

Temporarily replacing F by F_0, and h_j and h by h_{0j} and h_0, respectively, we would obtain:

$$\begin{aligned}
&\{1-F_0(t)\}h_{01}'(t)1_{S_1}(t) - F_0(t)h_{02}'(t)1_{S_2}(t) - f(t)\{\{1-F_0(t)\}g_1(t)1_{S_1}(t) + F_0(t)g_2(t)1_{S_2}(t)\}\\
&\qquad + F_0(t)\{1-F_0(t)\}\left\{\int_{u:(u,t)\in S}\frac{\partial_2 h_0(u,t)}{F_0(t)-F_0(u)}\,du - \int_{u:(t,u)\in S}\frac{\partial_1 h_0(t,u)}{F_0(u)-F_0(t)}\,du\right\}\\
&\qquad - f(t)F_0(t)\{1-F_0(t)\}\left\{\int_{u:(u,t)\in S}\frac{h_0(u,t)}{\{F_0(t)-F_0(u)\}^2}\,du + \int_{u:(t,u)\in S}\frac{h_0(t,u)}{\{F_0(u)-F_0(t)\}^2}\,du\right\} = 0.
\end{aligned}$$

that is:

$$\begin{aligned}
&f(t)\Biggl\{\{\{1-F_0(t)\}g_1(t)1_{S_1}(t) + F_0(t)g_2(t)1_{S_2}(t)\}\\
&\qquad\qquad + F_0(t)\{1-F_0(t)\}\left\{\int_{u:(u,t)\in S}\frac{h_0(u,t)}{\{F_0(t)-F_0(u)\}^2}\,du + \int_{u:(t,u)\in S}\frac{h_0(t,u)}{\{F_0(u)-F_0(t)\}^2}\,du\right\}\Biggr\}\\
&\quad = \{1-F_0(t)\}h_{01}'(t)1_{S_1}(t) - F_0(t)h_{02}'(t)1_{S_2}(t)\\
&\qquad\qquad + F_0(t)\{1-F_0(t)\}\left\{\int_{u:(u,t)\in S}\frac{\partial_2 h_0(u,t)}{F_0(t)-F_0(u)}\,du - \int_{u:(t,u)\in S}\frac{\partial_1 h_0(t,u)}{F_0(u)-F_0(t)}\,du\right\}.
\end{aligned}$$

This means that the coefficient of $f(t)$ in equation (4.11) stays away from zero if F belongs to a sufficiently small ball $B(F_0,\eta)$ around F_0, and $(h_1,h_2,h)\in U_\delta$ for small $\delta>0$. Denoting this coefficient by $c(t,h_1,h_2,h,F)$, we get the equation

$$\begin{aligned}
f(t) = \frac{1}{c(t,h_1,h_2,h,F)}&\Biggl\{\{1-F(t)\}h_1'(t) - F(t)h_2'(t)\\
&+F(t)\{1-F(t)\}\left\{\int_{u=0}^{t}\frac{\partial_2 h(u,t)}{F(t)-F(u)}\,du - \int_{u=t}^{M}\frac{\partial_1 h(t,u)}{F(u)-F(t)}\,du\right\}\Biggr\}.
\end{aligned}\tag{4.12}$$

The statement of the lemma now follows. □

The following lemma will be used to show that, with probability tending to one, $\tilde F_n$ belongs to the allowed class, for all large n, and is a consistent estimate of F_0.

Lemma 4.3. *Let, under the conditions of Lemma 4.1, $F^{(n)}\in B(F_0,\eta)$ be the solution of*

$$\phi(t;h_1^{(n)},h_2^{(n)},h^{(n)},F)=0,\,t\in[0,M],$$

where ϕ is defined by (4.5), and where $h^{(n)}\in C[S]$, $h_1^{(n)}\in C[S_1]$ an $h_2^{(n)}\in C[S_2]$ are non-negative functions which have bounded continuous derivatives on the supports S, S_1 and S_2, respectively, with finite limits approaching the boundary, respectively. Furthermore, let

$$\left\|h_j^{(n)}-h_{0j}\right\|\longrightarrow 0,\qquad \textit{and}\qquad \left\|\bigl(h_j^{(n)}\bigr)'-h_{0j}'\right\|\longrightarrow 0,\qquad j=1,2.$$

where, as before, $\|\cdot\|$ denotes the supremum norm on $C[S_j]$. Finally, let

$$\|h^{(n)}-h_0\|_S\longrightarrow 0\qquad \textit{and}\qquad \|\partial_j h^{(n)}-\partial_j h_0\|_S\longrightarrow 0,\qquad j=1,2, \tag{4.13}$$

where $\|\cdot\|_S$ is defined by (4.8). Then $F^{(n)}\to F_0$ in the supremum metric, as $n\to\infty$, and $F^{(n)}$ is strictly increasing on $[0,M]$ and satisfies $F^{(n)}(t)\in[0,1]$, $t\in[0,M]$, for all large n.

Proof. By Lemma 4.1 we have that $F^{(n)}$ tends to F_0 in the supremum norm on $C[0,M]$, since for any η we can choose a $\delta>0$ such that

$$\|F^{(n)}-F_0\|<\eta,$$

if $\|h_j^{(n)}-h_{0j}\|<\delta$ and $\|h^{(n)}-h_0\|<\delta$.

Using Lemma 4.2 we get that, under the conditions of the lemma, that $F^{(n)}$ is differentiable with a bounded derivative $f^{(n)}$, see (4.12). Specifically, (4.12) yields:

$$\begin{aligned} &f^{(n)}(t)\stackrel{\text{def}}{=}(F^{(n)})'(t)\\ &=\left\{F^{(n)}(t)\{1-F^{(n)}(t)\}\left\{\int_{u=0}^{t}\frac{\partial_2 h^{(n)}(u,t)}{F^{(n)}(t)-F^{(n)}(u)}\,du-\int_{u=t}^{M}\frac{\partial_1 h^{(n)}(t,u)}{F^{(n)}(u)-F^{(n)}(t)}\,du\right\}\right.\\ &\qquad\qquad\left.+\{1-F^{(n)}(t)\}\bigl(h_1^{(n)}\bigr)'(t)-F^{(n)}(t)\bigl(h_2^{(n)}\bigr)'(t)\right\}\Big/\,c(t,h_1^{(n)},h_2^{(n)},h^{(n)},F^{(n)}),\end{aligned}$$

where $c(t,h_1^{(n)},h_2^{(n)},h^{(n)},F^{(n)})$ stays away from zero, as $n\to\infty$. The corresponding density f_0 of the underlying model similarly has the representation

$$\begin{aligned} f_0(t)=&\left\{F_0(t)\{1-F_0(t)\}\left\{\int_{u:(u,t)\in S}\frac{\partial_2 h_0(u,t)}{F_0(t)-F_0(u)}\,du-\int_{u:(t,u)\in S}\frac{\partial_1 h_0(t,u)}{F_0(u)-F_0(t)}\,du\right\}\right.\\ &\qquad\qquad\left.+\{1-F_0(t)\}h_{01}'(t)-F_0(t)h_{02}'(t)\right\}\Big/\,c_0(t)\end{aligned} \tag{4.14}$$

where $c_0(t)$ is given by:

$$c_0(t)=g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t).$$

By

$$\|h_j^{(n)}-h_{0j}\|\to 0,\qquad \|h^{(n)}-h_0\|\to 0,\qquad \|F^{(n)}-F_0\|\to 0,$$

and (4.13), we now get:

$$\sup_{t\in[0,M]}\left|c\bigl(t,h_1^{(n)},h_2^{(n)},h^{(n)},F^{(n)}\bigr)-c_0(t)\right|\to 0.$$

Again using (4.13), we also get:

$$\|f^{(n)}-f_0\|\to 0,$$

that is: $f^{(n)}$ converges to f_0 in the supremum norm. Since f_0 stays away from zero on $[0,M]$, this means that $F^{(n)}$ is strictly increasing on $[0,M]$ for all sufficiently large n.

Furthermore, since $\phi(t,h_1^{(n)},h_2^{(n)},h^{(n)},F^{(n)})=0$, we get for large n, and t in a right neighborhood of 0,

$$F^{(n)}(t)=\frac{h_1^{(n)}(t)}{h_1^{(n)}(t)+h_2^{(n)}(t)+\{1-F^{(n)}(t)\}\int_{u=t}^M\frac{h^{(n)}(t,u)}{F^{(n)}(u)-F^{(n)}(t)}\,du}\ge 0,$$

since, by the convergence of $F^{(n)}$ to F_0, we may assume $1-F^{(n)}(t)>0$ for t in a neighborhood of 0, and since $h_j^{(n)}$ and $h^{(n)}$ are non-negative.

Likewise, if $F_0(M)=1$, we have, for t in a small left neighborhood of M,

$$\begin{aligned}&F^{(n)}(t)\{1-F^{(n)}(t)\}\int_{v=0}^t\frac{h^{(n)}(v,t)}{F^{(n)}(t)-F^{(n)}(v)}\,dv\\&=\left\{h_1^{(n)}(t)+h_2^{(n)}(t)\right\}F^{(n)}(t)-h_1^{(n)}(t)+F^{(n)}(t)\{1-F^{(n)}(t)\}\int_{v=t}^M\frac{h^{(n)}(v,t)}{F^{(n)}(t)-F^{(n)}(v)}\,dv\\&=h_2^{(n)}(t)F^{(n)}(t)-h_1^{(n)}(t)\{1-F^{(n)}(t)\},\end{aligned}$$

and hence, for t in a small left neighborhood of M,

$$\{1-F^{(n)}(t)\}\left\{F^{(n)}(t)\int_{v=0}^t\frac{h^{(n)}(v,t)}{F^{(n)}(t)-F^{(n)}(v)}\,dv+h_1^{(n)}(t)\right\}=h_2^{(n)}(t)F^{(n)}(t),$$

implying that, for all large n, $1-F^{(n)}(t)\ge 0$ for t in a neighborhood of M. This will a fortiori hold if $F_0(M)<1$. This shows that, for all large n and all $t\in[0,M]$, $F^{(n)}(t)\in[0,1]$. □

We still need to show that the estimates $\tilde h_{nj}$ of h_{0j} and $\tilde h_n$ of h_0 have the properties of h_j and h, as defined in Lemma 4.3.

LEMMA 4.4. *Let the conditions of Theorem 4.1 be satisfied and let the estimates $\tilde h_{nj}$ of and $\tilde h_n$ satisfy Condition 1.1. Then*

$$\left\|\tilde h_{nj}-h_{0j}\right\|\xrightarrow{p}0\qquad\text{and}\qquad\left\|\tilde h_{nj}'-h_{0j}'\right\|\xrightarrow{p}0\xrightarrow{p}0\qquad,\ j=1,2.$$

Moreover,

$$\|\tilde h_n-h_0\|_S\xrightarrow{p}0\qquad\text{and}\qquad\|\partial_j\tilde h_n-\partial_jh_0\|_S\xrightarrow{p}0,\qquad j=1,2.\tag{4.15}$$

Proof. Since we use boundary kernels near the boundary of $[0,M]$, $\tilde h_{nj}(t)$ is a consistent estimate of $h_{0j}(t)$ for each $t\in S_j$. For if $t\in[b_n,M-b_n]\cap S_j$ we just have:

$$E\tilde h_{n1}(t)=E\Delta_{11}K_{b_n}(t-T_1)=\int K_{b_n}(t-u)h_{01}(u)\,du=h_{01}(t)+O(n^{-2/5}),$$

where the remainder term is uniform in $t\in[b_n,M-b_n]\cap S_j$. Since $b_n\downarrow 0$, we have $b_n<\varepsilon$ for all large n, where ε is the "separation parameter" of Condition 1.1, and hence the boundary kernels are only relevant for $\tilde h_{n1}$ in a neighborhood of 0 and for $\tilde h_{n2}$ in a neighborhood of M.

If $t \in [0, b_n]$, we have:

$$\begin{aligned}
E\tilde h_{n1}(t) &= \alpha(t/b_n)E\Delta_{11}K_{b_n}(t-T_1) + \beta(t/b_n)E\left\{\Delta_{11}\frac{t-T_1}{b_n}K_{b_n}(t-T_1)\right\}\\
&= \alpha(t/b_n)\int_{u=0}^{M} K_{b_n}(t-u)h_{01}(u)\,du + \beta(t/b_n)\int_{u=0}^{M}\frac{t-u}{b_n}K_{b_n}(t-u)h_{01}(u)\,du\\
&= h_{01}(t)\int_{u=-1}^{t/b_n}\left\{\alpha(t/b_n)K(u) + \beta(t/b_n)K(u)\right\}\,du + O(n^{-2/5})\\
&= h_{01}(t) + O(n^{-2/5}),
\end{aligned}$$

again uniformly for $t \in [0, b_n]$. A similar computation can be made for $\tilde h_{n2}$ if $t \in [M - b_n, M]$. Since

$$\sup_{t\in S_1}\left|\tilde h_{n1}(t) - E\tilde h_{n1}(t)\right| = O_p\left(n^{-2/5}\sqrt{\log n}\right),$$

we now get the uniform convergence in probability of $\tilde h_{n1}$ to h_{01} on S_1, and similarly we have uniform convergence in probability of $\tilde h_{n2}$ to h_{02} on S_2.

Next we consider the derivative of $\tilde h_{n1}(t)$. If $t \in [b_n, M - b_n] \cap S_1$ we just have:

$$\begin{aligned}
&E\tilde h'_{n1}(t) = \frac{d}{dt}E\Delta_{11}K'_{b_n}(t-T_1) = \int \frac{d}{dt}K_{b_n}(t-u)h_{01}(u)\,du\\
&= b_n^{-1}\int K'(u)h_{01}(t-b_nu)\,du\\
&= b_n^{-1}\int K'(u)\left\{h_{01}(t) - b_nuh'_{01}(t) + \tfrac12 b_n^2u^2h'_{01}(t) - \tfrac16 b_n^3u^3h''_{01}(t)\right\}\,du + o\left(b_n^2\right)\\
&= h'_{01}(t) + O\left(b_n^2\right) = h'_{01}(t) + O\left(n^{-2/5}\right),
\end{aligned}$$

again uniformly in t. Since

$$\sup_{t\in[b_n,M-b_n]\cap S_1}\left|\tilde h'_{n1}(t) - E\tilde h'_{n1}(t)\right| = O_p\left(n^{-1/5}\sqrt{\log n}\right),$$

we only have to consider what happens near the boundary.

In treating the boundary kernels, we denote for simplicity b_n by b. If $t \in [0, b]$, we have

$$E\tilde h_{n1}(t) = \alpha\left(\frac{t}{b}\right)\int_{x=0}^{t+b}K_b(t-x)h_{01}(x)\,dx + \beta\left(\frac{t}{b}\right)\int_{x=0}^{t+b}\frac{t-x}{b}K_b(t-x)h_{01}(x)\,dx$$

This can also be written

$$E\tilde h_{n1}(t) = \int_{-1}^{t/b}\left\{\alpha\left(\frac{t}{b}\right)K(u) + \beta\left(\frac{t}{b}\right)uK(u)\right\}h_{01}(t-bu)\,du.$$

We write this in the form

$$E\tilde h_{n1}(t) = h_{01}(t) + \int_{-1}^{t/b}\left\{\alpha\left(\frac{t}{b}\right)K(u) + \beta\left(\frac{t}{b}\right)uK(u)\right\}\int_{t-bu}^{t}(w-t+bu)h''_{01}(w)\,dw\,du,$$

using a second order Taylor development of h_{01} with the integral remainder term. Hence

$$\begin{aligned}\frac{d}{dt}E\tilde{h}_{n1}(t) &= h_{01}'(t) + \frac{1}{b}\left\{\alpha\left(\frac{t}{b}\right)K\left(\frac{t}{b}\right) + \beta\left(\frac{t}{b}\right)\frac{t}{b}K\left(\frac{t}{b}\right)\right\}\int_0^t wh_{01}''(w)\,dw\\ &\qquad + \frac{1}{b}\int_{-1}^{t/b}\left\{\alpha'\left(\frac{t}{b}\right)K(u) + \beta'\left(\frac{t}{b}\right)uK(u)\right\}\int_{t-bu}^{t}(w-t+bu)h_{01}''(w)\,dw\,du\\ &\qquad - \int_{-1}^{t/b}\left\{\alpha\left(\frac{t}{b}\right)K(u) + \beta\left(\frac{t}{b}\right)uK(u)\right\}\int_{t-bu}^{t}h_{01}''(w)\,dw\,du\\ &\qquad + \int_{-1}^{t/b}\left\{\alpha\left(\frac{t}{b}\right)K(u) + \beta\left(\frac{t}{b}\right)uK(u)\right\}bu\,h_{01}''(t)\,du\\ &= h_{01}'(t) + O(b),\ b\downarrow 0.\end{aligned}$$

Note that, by Condition 1.2, the functions α, β, α' and β' are bounded on $[0,1]$. We also have:

$$\begin{aligned}\frac{d}{dt}E\tilde{h}_{n1}(t) &= \frac{1}{b}\left\{\alpha'\left(\frac{t}{b}\right)\int_{x=0}^{t+b}K_b(t-x)h_{01}(x)\,dx + \beta'\left(\frac{t}{b}\right)\int_{x=0}^{t+b}\frac{t-x}{b}K_b(t-x)h_{01}(x)\,dx\right\}\\ &\quad + \alpha\left(\frac{t}{b}\right)\int_{x=0}^{t+b}\frac{d}{dt}K_b(t-x)h_{01}(x)\,dx + \beta\left(\frac{t}{b}\right)\int_{x=0}^{t+b}\frac{t-x}{b}\frac{d}{dt}K_b'(t-x)h_{01}(x)\,dx\\ &\quad + \frac{1}{b}\beta\left(\frac{t}{b}\right)\int_{x=0}^{t+b}K_b(t-x)h_{01}(x)\,dx,\end{aligned}$$

so:

$$E\tilde{h}_{n1}'(t) = \frac{d}{dt}E\tilde{h}_{n1}(t) = h_{01}'(t) + O(b) = h_{01}'(t) + O\left(n^{-1/5}\right).$$

Since we have:

$$\sup_{t\in[0,b_n]}\left|\tilde{h}_{n1}'(t) - E\tilde{h}_{n1}'(t)\right| = O_p\left(n^{-1/5}\right),$$

we now also get that

$$\sup_{u\in S_1}\left|\tilde{h}_{n1}'(t) - h_{01}'(t)\right| = o_p(1).$$

The other cases can be treated in a similar way. □

We now get the following result.

Lemma 4.5. *Let the conditions of Theorem 4.1 be satisfied and let, for small $\eta > 0$, $F = \tilde{F}_n \in B(F_0,\eta)$ be the solution of the equation*

$$\phi(t;\tilde{h}_{n1},\tilde{h}_{n2},\tilde{h}_n,F) = 0,\ t\in[0,M].$$

where ϕ is defined by (4.5). Moreover, let $\|\cdot\|$ denote the supremum norm. Then:

(i) With probability tending to one, $\tilde{F}_n$ is strictly increasing on $[0,M]$, and satisfies $\tilde{F}_n(t)\in[0,1]$, $t\in[0,M]$, for all large n. Hence, with probability tending to one, $\tilde{F}_n$ coincides with the MSLE for large n and

$$\|\tilde{F}_n - F_0\| \xrightarrow{p} 0,\ n\to\infty.$$

(ii)

$$\|\tilde F_n - F_0\| = O_p\left(n^{-3/10}\right),\, n\to\infty.$$

(iii)

$$\|\tilde F_n - \bar F_n\| = O_p\left(n^{-3/5}\right),$$

where $\bar F_n$ is the solution in F of the linear integral equation

$$\begin{aligned}F(t)-F_0(t)+d_{F_0}(t)\Biggl\{\int_{u=0}^{t}\frac{g(u,t)\{F(t)-F_0(t)-F(u)+F_0(u)\}}{F_0(t)-F_0(u)}\,du\\ -\int_{u=t}^{M}\frac{g(t,u)\{F(u)-F_0(u)-F(t)+F_0(t)\}}{F_0(u)-F_0(t)}\,du\Biggr\}\\ =\frac{\tilde h_{n1}(t)\{1-F_0(t)\}-\tilde h_{n2}(t)F_0(t)}{\{1-F_0(t)\}g_1(t)+F_0(t)g_2(t)}\\ +d_{F_0}(t)\left\{\int_{u<t}\frac{\tilde h_n(u,t)}{F_0(t)-F_0(u)}\,du-\int_{u>v}\frac{\tilde h_n(t,u)}{F_0(u)-F_0(t)}\,du\right\},\end{aligned}\tag{4.16}$$

and where d_{F_0} is defined by

$$d_{F_0}(t)=\frac{F_0(t)\{1-F_0(t)\}}{g_1(t)\{1-F_0(t)+g_2(t)F_0(t)}\,.$$

Proof. Part (i) is an immediate consequence of Lemma 4.3.
(ii). We get, again using the approach of the implicit function theorem 10.2.1 in Banach spaces of [1], denoting the derivative w.r.t. (h_1,h_2,h) by D_1 and the derivative w.r.t. F by D_2:

$$\begin{aligned}\|\tilde F_n-F_0\| = \left\|\left[D_2\bar\phi(h_{01},h_{02},h_0,F_0)^{-1}\circ D_1\bar\phi(h_{01},h_{02},h_0,F_0)\right]\left(\tilde h_{n1}-h_{01},\tilde h_{n2}-h_{02},\tilde h_n-h_0\right)\right\|\\ +o_p\left(\|(\tilde h_{n1}-h_{01},\tilde h_{n2}-h_{02},\tilde h_n-h_0)\|\right),\end{aligned}\tag{4.17}$$

where the norm $\|\cdot\|$ on the left-hand side and the first norm on the right-hand side denote the supremum norm on $C[0,M]$ and the norm in the o_p-term denotes the norm

$$\|(h_1,h_2,h)\| = \max\left\{\|h_1\|,\|h_2\|,\|h\|_S\right\},$$

where the first two norms denote again the supremum norm and the third norm $\|\cdot\|_S$ is defined by (4.8).

By well-known results in density estimation, we have, if $b_n \asymp n^{-1/5}$,

$$\max\left(\|\tilde h_{n1}-h_{01}\|,\|\tilde h_{n2}-h_{02}\|\right)=O_p\left(n^{-2/5}\sqrt{\log n}\right),$$

The boundary kernels ensure that the rates are not spoiled by what happens at the boundary. So we have to determine the rate of convergence of $\|\tilde h_n-h_0\|_S$. We get:

$$\int_{u:(u,t)\in S}\left|\tilde h_n(u,t)-h_0(u,t)\right|\,du\le M^{1/2}\left\{\int_{u:(u,t)\in S}\left\{\tilde h_n(u,t)-h_0(u,t)\right\}^2\,du\right\}^{1/2},$$

and

$$E\int_{u:(u,t)\in S}\left\{\tilde h_n(u,t)-E\tilde h_n(u,t)\right\}^2\,du=O_p\left(\frac{1}{nb_n}\right)=O_p\left(n^{-4/5}\right).$$

For the bias we get, if $b_n<u<\gamma(u)+b_n\le t<M-b_n$, where γ is the function defining the lower boundary of the support S of g in Condition 1.1,

$$\begin{aligned}&E\tilde h_n(u,t)-h_0(u,t)=EK_{b_n}(u-T_1)K_{b_n}(t-U_1)\Delta_{12}\\&=\int\{F_0(u')-F_0(t')\}K_{b_n}(u-t')K_{b_n}(t-u')g(t',u')\,dt'\,du'-h_0(u,t)\\&=\int\{F_0(t-b_nv)-F_0(u-b_nw)\}K(v)K(w)g(u-b_nw,t-b_nv)\,dv\,dw-h_0(u,t)=O\left(b_n^2\right).\end{aligned}$$

If $\gamma(u)<t<\gamma(u)+b_n$ and $b_n<t<u<M-b_n$, we get:

$$E\tilde h_n(u,t)-h_0(u,t)=O(\gamma(u)-t).$$

The latter case gives a contribution of order $O(b_n^3)$ to the integral

$$\int_{u:(u,t)\in S}\left\{E\tilde h_n(u,t)-h_0(u,t)\right\}^2\,du.$$

The other cases can be treated in a similar way, with the conclusion that the dominating contribution to the integral

$$\left\{\int_{u:(u,t)\in S}\left\{\tilde h_n(u,t)-h_0(u,t)\right\}^2\,du\right\}^{1/2}$$

is coming from the bias near the lower boundary $\{(t,\gamma_n(t)):t\in[0,M]\}$ of S and is of order $O_p(n^{-3/10})$. So the conclusion is:

$$\|(\tilde h_{n1}-h_{01},\tilde h_{n2}-h_{02},\tilde h_n-h_0)\|=O_p\left(n^{-3/10}\right). \tag{4.18}$$

The derivative D_2 was computed in the proof of Lemma 4.1 (denoted by ∂_4 there) and the derivative D_1 is given by:

$$\begin{aligned}&\left[\left[D_1\bar\phi(h_{01},h_{02},h_0,F_0)\right](A)\right](t)\\&=B_1(t)\{1-F_0(t)\}-B_2(t)F_0(t)\\&\qquad+F_0(t)\{1-F_0(t)\}\left\{\int_{v=0}^t\frac{B(v,t)}{F_0(t)-F_0(v)}\,dv-\int_{u=t}^M\frac{B(t,u)}{F_0(u)-F_0(t)}\,du\right\},\end{aligned}$$

where B_1, B_2 and B are of the form

$$B_1=h_1-h_{01},\qquad B_2=h_2-h_{02},\qquad B=h-h_0.$$

Hence, defining $\bar F_n$ by

$$\bar F_n-F_0=-\left[D_2\bar\phi(h_{01},h_{02},h_0,F_0)^{-1}\circ D_1\bar\phi(h_{01},h_{02},h_0,F_0)\right](\tilde h_{n1}-h_{01},\tilde h_{n2}-h_{02},\tilde h_n-h_0), \tag{4.19}$$

we get that $F=\bar F_n$ is the solution of the linear integral equation

$$\left[D_2\bar\phi(h_{01},h_{02},h_0,F_0)\right]\left(F-F_0\right)=-\left[D_1\bar\phi(h_{01},h_{02},h_0,F_0)\right](\tilde h_{n1}-h_{01},\tilde h_{n2}-h_{02},\tilde h_n-h_0),$$

which, letting $A = F - F_0$ and $(B_1, B_2, B) = (\tilde h_{n1} - h_{01}, \tilde h_{n2} - h_{02}, \tilde h_n - h_0)$, boils down to the equation

$$\begin{aligned}
&\{h_{01}(t) + h_{02}(t)\}\,A(t) - \{1 - 2F_0(t)\}\left\{\int_{v=0}^{t} \frac{h_0(v,t)}{F_0(t) - F_0(v)}\,dv - \int_{u=t}^{M} \frac{h_0(t,u)}{F_0(u) - F_0(t)}\,du\right\}A(t)\\
&\qquad + F_0(t)\{1 - F_0(t)\}\left\{\int_{u=0}^{t} \frac{h_0(u,t)\{A(t) - A(u)\}}{\{F_0(t) - F_0(u)\}^2}\,du + \int_{u=t}^{M} \frac{h_0(t,u)\{A(t) - A(u)\}}{\{F_0(u) - F_0(t)\}^2}\,du\right\}\\
&= B_1(t)\{1 - F_0(t)\} - B_2(t)F_0(t)\\
&\qquad + F_0(t)\{1 - F_0(t)\}\left\{\int_{v=0}^{t} \frac{B(v,t)}{F_0(t) - F_0(v)}\,dv - \int_{u=t}^{M} \frac{B(t,u)}{F_0(u) - F_0(t)}\,du\right\},
\end{aligned}\tag{4.20}$$

We have:

$$\begin{aligned}
&\{h_{01}(t) + h_{02}(t)\}\,A(t) - \{1 - 2F_0(t)\}\left\{\int_{v=0}^{t} \frac{h_0(v,t)}{F_0(t) - F_0(v)}\,dv - \int_{u=t}^{M} \frac{h_0(t,u)}{F_0(u) - F_0(t)}\,du\right\}A(t)\\
&= \{g_1(t)F_0(t)(t) + g_2(t)\{1 - F_0(t)\}\}\,A(t) - \{1 - 2F_0(t)\}\{g_2(t) - g_1(t)\}A(t)\\
&= \{\{1 - F_0(t)\}g_1(t) + F_0(t)g_2(t)\}\,A(t).
\end{aligned}$$

Furthermore,

$$\begin{aligned}
&\int_{u=0}^{t} \frac{h_0(u,t)\{A(t) - A(u)\}}{\{F_0(t) - F_0(u)\}^2}\,du + \int_{u=t}^{M} \frac{h_0(t,u)\{A(t) - A(u)\}}{\{F_0(u) - F_0(t)\}^2}\,du\\
&= \int_{u=0}^{t} \frac{g(u,t)\{A(t) - A(u)\}}{F_0(t) - F_0(u)}\,du + \int_{u=t}^{M} \frac{g(t,u)\{A(t) - A(u)\}}{F_0(u) - F_0(t)}\,du.
\end{aligned}$$

Finally,

$$\begin{aligned}
&h_{01}(t)\{1 - F_0(t)\} - h_{02}(t)F_0(t)\\
&\qquad + F_0(t)\{1 - F_0(t)\}\left\{\int_{v=0}^{t} \frac{h_0(v,t)}{F_0(t) - F_0(v)}\,dv - \int_{u=t}^{M} \frac{h_0(t,u)}{F_0(u) - F_0(t)}\,du\right\} = 0.
\end{aligned}$$

So we obtain the linear integral equation (4.16) by dividing both sides of (4.20) by $\{1 - F_0(t)\}g_1(t) + F_0(t)g_2(t)$.

Hence $\bar F_n$ is the solution of the linear integral equation (4.16), and by (4.17) and (4.18),

$$\|\tilde F_n - F_0\| = \|\bar F_n - F_0\| + o_p\left(n^{-3/10}\right).$$

Note that we have:

$$\|\bar F_n - F_0\| = O_p\left(n^{-3/10}\right)$$

by the fact that $D_1\bar\phi(h_{01}, h_{02}, h_0, F_0)$ and $D_2\bar\phi(h_{01}, h_{02}, h_0, F_0)^{-1}$ are bounded linear mappings and hence:

$$\|\bar F_n - F_0\| = O_p\left(\|(\tilde h_{n1} - h_{01}, \tilde h_{n2} - h_{02}, \tilde h_n - h_0)\|\right) = O_p\left(n^{-3/10}\right).$$

(iii). The function $\tilde F_n$ satisfies the equation $\bar\phi(\tilde h_{n1}, \tilde h_{n2}, \tilde h_n, \tilde F_n) = 0$, where $\bar\phi$ is defined by (4.9). Hence:

$$\begin{aligned}
&\tilde h_{n1}(t)\{1 - \tilde F_n(t)\} - \tilde h_{n2}(t)\tilde F_n(t)\\
&\qquad + F(t)\{1 - F(t)\}\left\{\int_{v=0}^{t} \frac{\tilde h_n(v,t)}{\tilde F_n(t) - \tilde F_n(v)}\,dv - \int_{u=t}^{M} \frac{\tilde h_n(t,u)}{\tilde F_n(u) - \tilde F_n(t)}\,du\right\} = 0.
\end{aligned}$$

By (ii) we have $\|\tilde F_n - F_0\| = O_p(n^{-3/10})$, and hence:

$$\begin{aligned}
&\int_{v=0}^{t} \frac{\tilde h_n(v,t)}{\tilde F_n(t)-\tilde F_n(v)}\,dv\\
&\quad= \int_{v=0}^{t} \frac{\tilde h_n(v,t)}{F_0(t)-F_0(v)}\,dv - \int_{v=0}^{t} \frac{g(u,t)\{\tilde F_n(t)-F_0(t)-\tilde F_n(u)+F_0(u)\}}{F_0(t)-F_0(u)}\,du + O_p\left(n^{-3/5}\right),
\end{aligned}$$

and similarly

$$\begin{aligned}
&\int_{u=t}^{M} \frac{\tilde h_n(t,u)}{\tilde F_n(u)-\tilde F_n(t)}\,du\\
&\quad= \int_{u=t}^{M} \frac{\tilde h_n(t,u)}{F_0(u)-F_0(t)}\,du - \int_{u=t}^{M} \frac{g(t,u)\{\tilde F_n(u)-F_0(u)-\tilde F_n(t)+F_0(t)\}}{F_0(u)-F_0(t)}\,du + O_p\left(n^{-3/5}\right).
\end{aligned}$$

Hence we get:

$$\begin{aligned}
&\tilde F_n(t)-F_0(t)+d_{F_0}(t)\left\{\int_{u=0}^{t} \frac{g(u,t)\{\tilde F_n(t)-F_0(t)-\tilde F_n(u)+F_0(u)\}}{F_0(t)-F_0(u)}\,du\right.\\
&\qquad\qquad\left. - \int_{u=t}^{M} \frac{g(t,u)\{\tilde F_n(u)-F_0(u)-\tilde F_n(t)+F_0(t)\}}{F_0(u)-F_0(t)}\,du\right\}\\
&\quad= \frac{\tilde h_{n1}(t)\{1-F_0(t)\}-\tilde h_{n2}(t)F_0(t)}{\{1-F_0(t)\}g_1(t)+F_0(t)g_2(t)}\\
&\qquad + d_{F_0}(t)\left\{\int_{u=0}^{t} \frac{\tilde h_n(u,t)}{F_0(t)-F_0(u)}\,du - \int_{u=t}^{M} \frac{\tilde h_n(t,u)}{F_0(u)-F_0(t)}\,du\right\} + O_p\left(n^{-3/5}\right),
\end{aligned}\tag{4.21}$$

uniformly for $t \in [0,M]$, implying

$$\begin{aligned}
\tilde F_n &= -\left[D_2\bar\phi(h_{01},h_{02},h_0,F_0)^{-1}\circ D_1\bar\phi(h_{01},h_{02},h_0,F_0)\right]\left(\tilde h_{n1}-h_{01},\tilde h_{n2}-h_{02},\tilde h_n-h_0\right)\\
&\qquad\qquad + O_p\left(n^{-3/5}\right)\\
&= \bar F_n + O_p\left(n^{-3/5}\right).
\end{aligned}$$

□

We now have in principle solved our problem, since we have shown that $\tilde F_n$ is locally asymptotically equivalent to the solution $\bar F_n$ of a linear integral equation. Since $\tilde F_n$ coincides with the MSLE for large n, the MSLE is also locally asymptotically equivalent with $\bar F_n$. However, to get an explicit expression for the bias and variance of the MSLE, we now study a still simpler "toy estimator", which turns out also to be locally asymptotically equivalent to the MSLE.

LEMMA 4.6. *Let the toy estimator $F = F_n^{toy}$ be defined as the solution of the equation*

$$\begin{aligned}
&\{F(t)-F_0(t)\}\left\{1+d_{F_0}(t)\left\{\int_{u<t}\frac{g(u,t)}{F_0(t)-F_0(u)}\,dt+\int_{u>v}\frac{g(t,u)}{F_0(u)-F_0(t)}\,du\right\}\right\}\\
&\quad= \frac{\tilde h_{n1}(t)\{1-F_0(t)\}-\tilde h_{n2}(t)F_0(t)}{\{1-F_0(t)\}g_1(t)+F_0(t)g_2(t)}\\
&\qquad + d_{F_0}(t)\left\{\int_{u<t}\frac{\tilde h_n(u,t)}{F_0(t)-F_0(u)}\,du-\int_{u>v}\frac{\tilde h_n(t,u)}{F_0(u)-F_0(t)}\,du\right\}.
\end{aligned}\tag{4.22}$$

Then, under the conditions of Theorem 4.1,

$$\sqrt{nb_n}\left\{F_n^{toy}(v)-F_0(v)-\frac{\beta(v)b_n^2}{2\sigma_1(v)}\right\}\xrightarrow{\mathcal{D}} N\left(0,\sigma(v)^2\right),$$

where $b_n \asymp n^{-1/5}$ *and* $\beta(v)$, $\sigma_1(v)$ *and* $\sigma(v)$ *are defined as in Theorem 4.1.*

Remark 4.1. In Lemma 4.6 a toy estimator is introduced, which plays a similar role as the toy estimator in the study of the ordinary MLE for interval censoring, introduced in [5] and [11] (the term "toy estimator" was coined by Jon Wellner). It is called a toy estimator because we cannot use it in an actual sample, since F_0 is unknown (and is in fact the object we want to estimate). Actually, the solution $\bar F_n$ of the linear integral equation (4.16) in part (iii) of Lemma 4.5 is also a toy estimator in this sense (but does not produce explicit expressions for the expectation and variance of the asymptotic distribution).

Proof of Lemma 4.6. We have:

$$\tilde h_{n1}(v)=\frac1n\sum_{i=1}^n K_{b_n}(v-T_i)\Delta_{i1}$$

and hence

$$\begin{aligned}\mathrm{var}\left(\tilde h_{n1}(v)\right)&=\frac1n\mathrm{var}\left(K_{b_n}(v-T_1)\Delta_{11}\right)\\&=\frac1n EK_{b_n}(v-T_1)^2\bigl(\Delta_{11}-F_0(T_1)\bigr)^2\sim\frac{F_0(v)\{1-F_0(v)\}g_1(v)}{nb_n}\int K(u)^2\,du.\end{aligned}$$

Likewise

$$\mathrm{var}\left(\tilde h_{n2}(v)\right)\sim\frac{F_0(v)\{1-F_0(v)\}g_2(v)}{nb_n}\int K(u)^2\,du.$$

Furthermore,

$$\begin{aligned}\mathrm{covar}\left(\tilde h_{n1}(v),\tilde h_{n2}(v)\right)&=\frac1n EK_{b_n}(v-T_1)K_{b_n}(v-U_1)\bigl(\Delta_{11}-F_0(T_1)\bigr)\bigl(\Delta_{13}-F_0(U_1)\bigr)\\&=-\frac1n EK_{b_n}(v-T_1)K_{b_n}(v-U_1)F_0(T_1)F_0(U_1)\sim-\frac{F_0(v)^2g(v,v)}{nb_n}\int K(u)^2\,du=0,\end{aligned}$$

using the "separation condition" $g(v,v)=0$. So we obtain:

$$\mathrm{var}\left(\frac{\{1-F_0(v)\}\tilde h_{n1}(v)-F_0(v)\tilde h_{n2}(v)}{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)}\right)\sim\frac{F_0(v)\{1-F_0(v)\}\left(\{1-F_0(v)\}^2g_1(v)+F_0(v)^2g_2(v)\right)}{nb_n\bigl\{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)\bigr\}^2}.$$

Furthermore,

$$\int_{t<v}\frac{\tilde h_n(t,v)}{F_0(v)-F_0(t)}\,dt=n^{-1}\sum_{i=1}^n K_{b_n}(v-U_i)\Delta_{i2}\int_{t<v}\frac{K_{b_n}(t-T_i)}{F_0(v)-F_0(t)}\,dt,$$

and hence

$$
\begin{aligned}
&\mathrm{var}\left(\int_{t<v}\frac{\tilde h_n(t,v)}{F_0(v)-F_0(t)}\,dt\right)=n^{-1}\mathrm{var}\left(K_{b_n}(v-U_1)\Delta_{12}\int_{t<v}\frac{K_{b_n}(t-T_1)}{F_0(v)-F_0(t)}\,dt\right)\\
&=n^{-1}EK_{b_n}(v-U_1)^2\bigl(\Delta_{12}-F_0(U_1)+F_0(T_1)\bigr)^2\left\{\int_{t<v}\frac{K_{b_n}(t-T_1)}{F_0(v)-F_0(t)}\,dt\right)^2\\
&=n^{-1}EK_{b_n}(v-U_1)^2\left\{F_0(U_1)-F_0(T_1)\right\}\left\{1-F_0(U_1)+F_0(T_1)\right\}\left\{\int_{t<v}\frac{K_{b_n}(t-T_1)}{F_0(v)-F_0(t)}\,dt\right)^2\\
&\sim\frac{1}{nb_n}\int_{t<v}\frac{1-F_0(v)+F_0(t)}{F_0(v)-F_0(t)}\,g(t,v)\,dt\int K(u)^2\,du.
\end{aligned}
$$

Likewise

$$
\mathrm{var}\left(\int_{w>v}\frac{\tilde h_n(v,w)}{F_0(w)-F_0(v)}\,dw\right)\sim\frac{1}{nb_n}\int_{w>v}\frac{1-F_0(w)+F_0(v)}{F_0(w)-F_0(v)}\,g(v,w)\,dw\int K(u)^2\,du.
$$

Finally,

$$
\begin{aligned}
&\mathrm{covar}\left(\tilde h_{n2}(v),\int_{t<v}\frac{\tilde h_n(t,v)}{F_0(v)-F_0(t)}\,dt\right)\\
&=\frac1n EK_{b_n}(v-U_1)^2\bigl(\Delta_{13}-\{1-F_0(U_1)\}\bigr)\left(\Delta_{12}-F_0(U_1)+F_0(T_1)\right)\int_{t<v}\frac{K_{b_n}(t-T_1)}{F_0(v)-F_0(t)}\,dt\\
&=-\frac1n EK_{b_n}(v-U_1)^2\{1-F_0(U_1)\}\{F_0(U_1)-F_0(T_1)\}\int_{t<v}\frac{K_{b_n}(t-T_1)}{F_0(v)-F_0(t)}\,dt\\
&\sim-\frac{\{1-F_0(v)\}g_2(v)}{nb_n}\int K(u)^2\,du.
\end{aligned}
$$

and similarly:

$$
\mathrm{covar}\left(\tilde h_{n1}(v),\int_{w>v}\frac{\tilde h_n(v,w)}{F_0(w)-F_0(v)}\,dw\right)\sim-\frac{F_0(v)g_1(v)}{nb_n}\int K(u)^2\,du.
$$

Combining these facts, we obtain that the variance of the right-hand side of (4.22) is given by:

$$
\begin{aligned}
&\frac{F_0(v)\{1-F_0(v)\}\left(\{1-F_0(v)\}^2g_1(v)+F_0(v)^2g_2(v)\right)}{nb_n\bigl\{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)\bigr\}^2}\int K(u)^2\,du\\
&\qquad+\frac{d_{F_0}(v)^2}{nb_n}\left\{\int_{t<v}\frac{g(t,v)}{F_0(v)-F_0(t)}\,dt+\int_{w>v}\frac{g(v,w)}{F_0(w)-F_0(v)}\,dw-g_1(v)-g_2(v)\right\}\int K(u)^2\,du\\
&\qquad+\frac{2d_{F_0}(v)^2}{nb_n}\left\{g_1(v)+g_2(v)\right\}\int K(u)^2\,du\\
&=\frac{F_0(v)\{1-F_0(v)\}\left(\{1-F_0(v)\}^2g_1(v)+F_0(v)^2g_2(v)\right)}{nb_n\bigl\{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)\bigr\}^2}\int K(u)^2\,du\\
&\qquad+\frac{d_{F_0}(v)^2}{nb_n}\left\{\int_{t<v}\frac{g(t,v)}{F_0(v)-F_0(t)}\,dt+\int_{w>v}\frac{g(v,w)}{F_0(w)-F_0(v)}\,dw+g_1(v)+g_2(v)\right\}\int K(u)^2\,du\\
&=\frac{d_{F_0}(v)}{nb_n}\left\{1+d_{F_0}(v)\left\{\int_{t<v}\frac{g(t,v)}{F_0(v)-F_0(t)}\,dt+\int_{w>v}\frac{g(v,w)}{F_0(w)-F_0(v)}\,dw\right\}\right\}\int K(u)^2\,du.
\end{aligned}
$$

Hence we get that the asymptotic variance at a fixed interior point v of the solution F of the equation (4.22) is given by:

$$\frac{d_{F_0}(v)}{nb_n\sigma_1}\int K(u)^2\,du,$$

where σ_1 is defined by (4.2).

We still have to compute the bias. We have:

$$\begin{aligned}
&\frac{\{1-F_0(v)\}\bigl\{\tilde h_1(v)-h_1(v)\bigr\}-F_0(v)\bigl\{\tilde h_2(v)-h_2(v)\bigr\}}{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)}\\
&\qquad+d_{F_0}(v)\int_{t<v}\frac{\tilde h(t,v)-h(t,v)}{F_0(v)-F_0(t)}\,dt-d_{F_0}(v)\int_{u>v}\frac{\tilde h_n(v,u)-h(v,u)}{F_0(u)-F_0(v)}\,du\\
&=\frac{\{1-F_0(v)\}h_1''(v)-F_0(v)h_2''(v)}{2\,\{g_1(v)\{1-F_0(v)\}+F_0(v)g_2(v)\}}b_n^2\int u^2K(u)\,du\\
&\qquad+\tfrac12 b_n^2 d_{F_0}(v)\left\{\int_{t<v}\frac{\frac{\partial^2}{\partial v^2}h(t,v)}{F_0(v)-F_0(t)}\,dt-\int_{u>v}\frac{\frac{\partial^2}{\partial v^2}h(v,u)}{F_0(u)-F_0(v)}\,du\right\}\int u^2K(u)\,du,
\end{aligned}$$

where

$$\tilde h_1(t)=\int K_{b_n}(t-u)F_0(u)\,g_1(u)\,du,\qquad \tilde h_2(t)=\int K_{b_n}(t-u)\{1-F_0(u)\}\,g_2(u)\,du \tag{4.23}$$

and

$$\tilde h(t,u)=\int K_{b_n}(t-v)K_{b_n}(u-w)\{F_0(w)-F_0(v)\}\,g(v,w))\,dv\,dw. \tag{4.24}$$

Moreover,

$$\begin{aligned}
&\{1-F_0(v)\}h_1(v)-F_0(v)h_2(v)\\
&\qquad+F_0(v)\{1-F_0(v)\}\left\{\int_{t<v}\frac{h(t,v)}{F_0(v)-F_0(t)}\,dt-\int_{u>v}\frac{h(v,u)}{F_0(u)-F_0(v)}\,du\right\}\\
&=F_0(v)\{1-F_0(v)\}\,\{g_1(v)-g_2(v)\}-F_0(v)\{1-F_0(v)\}\,\{g_1(v)-g_2(v)\}=0.
\end{aligned}$$

This yields the result of the lemma, since we can use the central limit theorem for i.i.d. random variables on the right-hand side of (4.22), using

$$\begin{aligned}
&\frac{\tilde h_{n1}(t)\{1-F_0(t)\}-\tilde h_{n2}(t)F_0(t)}{\{1-F_0(t)\}g_1(t)+F_0(t)g_2(t)}\\
&\qquad+d_{F_0}(t)\left\{\int_{u<t}\frac{\tilde h_n(u,t)}{F_0(t)-F_0(u)}\,du-\int_{u>t}\frac{\tilde h_n(t,u)}{F_0(u)-F_0(t)}\,du\right\}\\
&=n^{-1}\sum_{i=1}^n\left\{\frac{\{1-F_0(t)\}K_{b_n}(t-T_i)\Delta_{i1}-F_0(t)K_{b_n}(t-U_i)\Delta_{i3}}{\{1-F_0(t)\}g_1(t)+F_0(t)g_2(t)}\right.\\
&\qquad+d_{F_0}(t)K_{b_n}(t-U_i)\Delta_{i2}\int_{u<t}\frac{K_{b_n}(u-T_i)}{F_0(t)-F_0(u)}\,du\\
&\qquad\left.-d_{F_0}(t)K_{b_n}(t-T_i)\Delta_{i2}\int_{u>t}\frac{K_{b_n}(u-U_i)}{F_0(u)-F_0(t)}\,du\right\}.
\end{aligned} \tag{4.25}$$

□

We now show that the solution of the linear integral equation (4.16) is equivalent in first order to the toy estimator of Lemma 4.6.

Lemma 4.7. *Let, under the conditions of Theorem 4.1, F_n^{toy} solve equation (4.22) of Lemma 4.6 and let $\bar F_n$ be the solution of the linear integral equation (4.16). Then:*

$$\bar F_n(t) = F_n^{toy}(t) + O_p\left(n^{-1/2}\right).$$

Proof. It is clear from the definition of $\bar F_n$ that we only have to show:

$$\int_{u<t} \frac{\tilde h_n(u,t)\{\bar F_n(u) - F_0(u)\}}{\{F_0(t)-F_0(u)\}^2}\,du = O_p\left(n^{-1/2}\right),$$

and

$$\int_{u>t} \frac{\tilde h_n(t,u)\{\bar F_n(u) - F_0(u)\}}{\{F_0(u)-F_0(t)\}^2}\,du = O_p\left(n^{-1/2}\right).$$

We only prove the first relation, since the proof of the second relation is completely similar.

Defining $A_n = \bar F_n - F_0$, $B_{n1} = \tilde h_{n1} - h_{10}$, $B_{n2} = \tilde h_{n2} - h_{10}$ and $B_n = \tilde h_n - h_0$, we have, for a bounded weight function w,

$$\begin{aligned}
&\int_{t=0}^{x} w(t,x)\left\{g_1(t)\{1-F_0(t)\} + g_2(t)F_0(t)\right\} A_n(t)\,dt\\
&\qquad + \int_{t=0}^{x} w(t,x)F_0(t)\{1-F_0(t)\}\left\{\int_{u=0}^{t} \frac{h_0(u,t)\{A_n(t)-A_n(u)\}}{\{F_0(t)-F_0(u)\}^2}\,du\right.\\
&\qquad\qquad\qquad\qquad\qquad\qquad\left. + \int_{u=t}^{M} \frac{h_0(t,u)\{A_n(u)-A_n(t)\}}{\{F_0(u)-F_0(t)\}^2}\,du\right\} dt\\
&= \int_{t=0}^{x} w(t,x)\left\{B_{n1}(t)\{1-F_0(t)\} - B_{n2}(t)F_0(t)\right\} dt\\
&\qquad + \int_{t=0}^{x} w(t,x)F_0(t)\{1-F_0(t)\}\left\{\int_{v=0}^{t} \frac{B_n(v,t)}{F_0(t)-F_0(v)}\,dv - \int_{u=t}^{M}\frac{B_n(t,u)}{F_0(u)-F_0(t)}\,du\right\} dt.
\end{aligned} \tag{4.26}$$

The right-hand side of this equation is clearly $O_p(n^{-1/2})$, uniformly in $x \in [0,M]$, where we use the fact that the kernel estimators also behave well at the boundary, since we use boundary kernels.

Since we can write, for example,

$$\begin{aligned}
&\int_{t=0}^{x} w(t,x)F_0(t)\{1-F_0(t)\}\int_{u=0}^{t} \frac{h_0(u,t)A_n(u)}{\{F_0(t)-F_0(u)\}^2}\,du\,dt\\
&= \int_{t=0}^{x}\left\{\int_{y=t}^{x} w(y,x)F_0(y)\{1-F_0(y)\}\frac{h_0(t,y)}{\{F_0(y)-F_0(t)\}^2}\,du\right\} A_n(t)\,dt,
\end{aligned}$$

and

$$\begin{aligned}
&\int_{t=0}^{x} w(t,x)F_0(t)\{1-F_0(t)\}\int_{u=t}^{M} \frac{h_0(t,u)A_n(u)}{\{F_0(u)-F_0(t)\}^2}\,du\\
&= \int_{t=0}^{x}\left\{\int_{u=0}^{t} w(u,x)F_0(u)\{1-F_0(u)\}\frac{h_0(u,t)}{\{F_0(t)-F_0(u)\}^2}\,du\right\} A_n(t)\,dt\\
&\qquad\qquad + \int_{t=x}^{M}\left\{\int_{u=0}^{x} w(u,x)F_0(u)\{1-F_0(u)\}\frac{h_0(u,t)}{\{F_0(t)-F_0(u)\}^2}\,du\right\} A_n(t)\,dt
\end{aligned}$$

the left-hand side of (4.26) is of the form

$$\int_0^x \bar w(t,x)A_n(t)\,dt,$$

for a bounded weight function $\bar w$, depending on w, F_0, h_{0j} and h_0. Defining

$$\begin{aligned}a(t) &\overset{\text{def}}{=} g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t)\\ &\qquad + F_0(t)\{1-F_0(t)\}\left\{\int_{u=0}^t \frac{h_0(u,t)}{\{F_0(t)-F_0(u)\}^2}\,du+\int_{u=t}^M \frac{h_0(t,u)}{\{F_0(u)-F_0(t)\}^2}\,du\right\}\,dt\end{aligned}$$

which stays away from zero, for $t\in[0,M]$, we can solve the (Fredholm) integral equation

$$\begin{aligned}a(x)w(t,x) &= \frac{h_0(t,x)}{\{F_0(x)-F_0(t)\}^2}+\int_{y=t}^x w(y,x)F_0(y)\{1-F_0(y)\}\frac{h_0(t,y)}{\{F_0(y)-F_0(t)\}^2}\,dy\\ &\qquad+\int_{y=0}^{t\wedge x} w(y,x)F_0(y)\{1-F_0(y)\}\frac{h_0(y,t)}{\{F_0(t)-F_0(y)\}^2}\,dy \end{aligned}\tag{4.27}$$

for the function $t\mapsto w(t,x)$, $t\in[0,x]$, and next use (4.27) to define $w(t,x)$ for $t\in(x,M]$.

To show that the solution exists, we write, for $t\le x$, the equation in the form

$$\begin{aligned}&\left\{g_1(t)\{1-F_0(t)\}+g_2(t)F_0(t)+F_0(t)\{1-F_0(t)\}\int_{u=x}^M \frac{h_0(t,u)}{\{F_0(u)-F_0(t)\}^2}\,du\,dt\right\}w(t,x)\\ &=\frac{h_0(t,x)}{\{F_0(x)-F_0(t)\}^2}\\ &\qquad+\int_{y=t}^x \frac{h_0(t,y)}{\{F_0(y)-F_0(t)\}^2}\{F_0(y)\{1-F_0(y)\}w(y,x)-F_0(t)\{1-F_0(t)\}w(t,x)\}\,dy\\ &\qquad\quad-\int_{y=0}^t \frac{h_0(y,t)}{\{F_0(t)-F_0(y)\}^2}\ \{F_0(t)\{1-F_0(t)\}w(t,x)-F_0(y)\{1-F_0(y)\}w(y,x)\}\,dy,\end{aligned}$$

and use similar techniques as in the treatment of the integral equation

$$\left[\partial_4\bar\phi(h_{01},h_{02},h_0,F_0)\right](A)=0,$$

in the proof of Lemma 4.1 to show that the homogeneous equation only has the trivial solution zero, which means by the Fredholm theory that there exists a (unique) solution.

For this choice of the weight function w we find:

$$\begin{aligned}&\int_{u=0}^t \frac{h_0(u,t)A_n(u)}{\{F_0(t)-F_0(u)\}^2}\,du\\ &=\int_{u=0}^t w(u,t)\,\{g_1(u)\{1-F_0(u)\}+g_2(u)F_0(u)\}\,A_n(u)\,du\\ &\qquad+\int_{u=0}^t w(u,t)F_0(u)\{1-F_0(u)\}\left\{\int_{v=0}^u \frac{h_0(v,u)\{A_n(u)-A_n(v)\}}{\{F_0(u)-F_0(v)\}^2}\,dv\right.\\ &\qquad\qquad\left.+\int_{v=u}^M \frac{h_0(u,v)\{A_n(v)-A_n(u)\}}{\{F_0(v)-F_0(u)\}^2}\,dv\right\}\,du\\ &=\int_{u=0}^t w(u,t)\,\{B_{n1}(u)\{1-F_0(u)\}-B_{n2}(u)F_0(u)\}\,du\\ &\qquad+\int_{u=0}^t w(u,t)F_0(u)\{1-F_0(u)\}\left\{\int_{v=0}^u \frac{B_n(v,u)}{F_0(u)-F_0(v)}\,dv-\int_{v=u}^M \frac{B_n(u,v)}{F_0(v)-F_0(u)}\,dv\right\}\,du,\end{aligned}$$

and the expression on the right-hand side is $O_p\left(n^{-1/2}\right)$, uniformly for $t \in [0, M]$.

The term

$$\int_t^M \frac{h_0(t,u)A_n(u)}{\{F_0(u)-F_0(t)\}^2}\,du$$

can be treated in a similar way. The conclusion of the lemma now follows. ☐

Theorem 4.1 now follows from the asymptotic equivalence of the the toy estimator with the solution $\bar{F}_n$ of the linear integral equation (4.16) and the asymptotic equivalence of $\bar{F}_n$ with the MSLE.

5. Concluding remarks and open problems.

In the preceding it was shown that, under the so-called separation hypothesis, the MSLE locally converges to the underlying distribution function at rate $n^{-2/5}$, if we use bandwidths $b_n \asymp n^{-1/5}$ in the estimates $\tilde{h}_{nj}$ and $\tilde{h}_n$. The asymptotic (normal) distribution was also determined. The results can be used to construct a two-sample likelihood ratio test, of the same type as the test, discussed in [7] for the current status model, but this is not done in the present paper because of reasons of space.

If the separation hypothesis does not hold, which means that we can have arbitrarily small observation intervals, the asymptotic behavior of the MSLE is still unknown. In this situation the local asymptotic limit distribution for the ordinary MLE is also still unknown, although it is conjectured that the rate $n^{-1/3}$, holding under the separation hypothesis, is improved to the rate $(n\log n)^{-1/3}$ in this case. There even exists a conjectured limit distribution in this case, put forward in [5] (see also [11]). Supporting evidence for this conjecture is given in a simulation study in [10], but a proof is still missing. The latter paper also gives simulation results for the SMLE, and it is conjectured that the SMLE will locally have a similar behavior as the MSLE if the separation hypothesis hold. The asymptotic behavior of the SMLE again has to be deduced from an associated integral equation. This matter is further discussed in [10].

References.

[1] Jean Dieudonné, *Foundations of modern analysis*, Academic Press, New York, 1969, Enlarged and corrected printing, Pure and Applied Mathematics, Vol. 10-I. MR0349288 (50 #1782)

[2] Ronald B. Geskus and Piet Groeneboom, *Asymptotically optimal estimation of smooth functionals for interval censoring. I*, Statist. Neerlandica **50** (1996), no. 1, 69–88. MR1381209 (97k:62039)

[3] ———, *Asymptotically optimal estimation of smooth functionals for interval censoring. II*, Statist. Neerlandica **51** (1997), no. 2, 201–219. MR1466426 (99d:62015)

[4] ———, *Asymptotically optimal estimation of smooth functionals for interval censoring, case* 2, Ann. Statist. **27** (1999), no. 2, 627–674. MR1714713 (2000j:60044)

[5] Piet Groeneboom, *Nonparametric maximum likelihood estimators for interval censoring and deconvolution*, Technical Report 378, Department of Statistics, Stanford University, 1991.

[6] ———, *Lectures on inverse problems*, Lectures on probability theory and statistics (Saint-Flour, 1994), Lecture Notes in Math., vol. 1648, Springer, Berlin, 1996, pp. 67–164. MR1600884 (99c:62092)

[7] ———, *Likelihood ratio type two-sample tests for current status data*, To appear in the Scandinavian Journal of Statistics, 2012.

[8] Piet Groeneboom, Geurt Jongbloed, and Birgit I. Witte, *Maximum smoothed likelihood estimation and smoothed maximum likelihood estimation in the current status model*, Ann. Statist. **38** (2010), no. 1, 352–387. MR2589325 (2011c:62098)

[9] ———, *A maximum smoothed likelihood estimator in the current status continuous mark model*, J. Nonparametr. Stat. **24** (2012), no. 1, 85–101.

[10] Piet Groeneboom and Tom Ketelaars, *Estimators for the interval censoring problem*, Electron. J. Stat. **5** (2011), 1797–1845.

[11] Piet Groeneboom and Jon A. Wellner, *Information bounds and nonparametric maximum likelihood estimation*, DMV Seminar, vol. 19, Birkhäuser Verlag, Basel, 1992. MR1180321 (94k:62056)

[12] Geurt Jongbloed, *The iterative convex minorant algorithm for nonparametric estimation*, J. Comput. Graph. Statist. **7** (1998), no. 3, 310–321. MR1646718

[13] Marloes H. Maathuis and Jon A. Wellner, *Inconsistency of the MLE for the joint distribution of interval-censored survival times and continuous marks*, Scand. J. Statist. **35** (2008), no. 1, 83–103. MR2391560 (2009e:62392)

[14] Wei-Yann Tsai, Sue Leurgans, and John Crowley, *Nonparametric estimation of a bivariate survival function in the presence of censoring*, Ann. Statist. **14** (1986), no. 4, 1351–1365. MR868304 (88a:62108)

Delft Institute of Applied Mathematics, Delft University of Technology,
Mekelweg 4, 2628 CD Delft, The Netherlands
E-mail: P.Groeneboom@tudelft.nl, URL: http://dutiosc.twi.tudelft.nl/˜pietg/